The Third National Workshop on the SKA Project - The Italian Route to the SKAO Revolution



Contribution ID: 38

Type: not specified

## The MeerKAT Fornax Survey: Rapid HI removal in dwarf galaxies

Thursday, 7 October 2021 11:20 (20 minutes)

I will present some of the first results from the MeerKAT Fornax Survey (MFS), that show that dwarf galaxies rapidly lose their cool gas once they have entered into the cluster. The MFS is now 25% complete, and thanks to the exquisite sensitivity and resolution of the MeerKAT telescope, we are exploring a new observational parameter space and have detected HI in a population of dwarf (Mr > -18.5) galaxies down to a HI mass of 10<sup>6</sup> Msol. Only 14 out of 300 dwarfs in the current mosaic are detected in HI, showing that the majority of dwarfs in the Fornax cluster have already lost their HI. For the dwarfs with HI, a dichotomy exists - either the dwarf galaxies in between (i.e. transitioning from HI-normal to HI-poor) these two HI mass-to-luminosity ratios, this suggests that a dwarf galaxy with HI will very rapidly lose it's HI, even before the first pericentric passage. We also find that the majority of dwarfs with HI are at least 80 kpc away from the closest, massive galaxy. This may be evidence that pre-processing assists in the removal of HI from dwarf satellites, accelerating the loss of HI while infalling to and within the cluster.

## **Reasearch** area

HI galaxy science

Primary author: KLEINER, Dane (Istituto Nazionale di Astrofisica (INAF))
Co-author: THE MFS TEAM
Presenter: KLEINER, Dane (Istituto Nazionale di Astrofisica (INAF))
Session Classification: Nearby Universe & Galaxy Evolution