



# ASTRI Mini-Array Core Science

Stefano Vercellone – INAF Osservatorio Astronomico di Brera  
for the ASTRI Project

**OAS VHE Meeting, 08-09.06.2022**



See also Talk by S. Scuderi

## ASTRI-Horn Prototype

INAF-led Project funded by Italian Ministry of Research

End-to-end prototype installed and operational on Mount Etna volcano (Sicily, Italy)

**First detection of a gamma-ray source** (Crab Nebula) above  $5\sigma$  **with a dual-mirror, Schwarzschild-Couder Chrenkov telescope** (Lombardi et al., 2020)



## Array of 9 ASTRI telescopes

INAF-led Project with international partners: Univ. of Sao Paulo/FPESP (Brazil), North-West Univ. (S. Africa), IAC (Spain), FGG, ASI/SSDC, Univ. of Padova, Perugia and INFN

**Being deployed at the *Observatorio del Teide*** (Spain) in collaboration with IAC and FGG-INAF.

**First 4 yr → *Core Science***, following 4 yr → *Observatory Science*. **Science operation → 2024**



# Simulation and Data Processing Systems

## Simulation System


Software System that provides (low-level) scientific simulated data for the development of reconstruction algorithms, the assessment of the system performance, and the characterization of real observations

## Data Processing System

Software System used to perform the offline software stereo trigger, to calibrate, reconstruct, select, and analyze the ASTRI Mini-Array Cherenkov and SI<sup>3</sup> data, and to check the data quality


Main components:

- Stereo Event Builder
- Cherenkov Data Pipeline
- Intensity Interferometry Data Pipeline
- Calibration Software



# Cherenkov Data Pipeline

SPIE 991315 (2016); SPIE 107070 (2018)



Functional design layout of the ASTRI data reduction and scientific analysis software package (**A-SciSoft**). From left to right, functional breakdown stages are shown as dotted boxes, I/O data as grey cylinders (black: main pipeline I/O data; grey: auxiliary I/O data), basic software functionalities as dashed boxes, data levels as grey boxes (thin: telescope-wise data; thick: array-wise data), and data processing executables as solid boxes. The external packages for the scientific analysis, i.e. `ctools` and `Gammapy`, are written in bright grey.

## A-SciSoft (ASTRI Scientific Software, aka **astripipe**)

- End-to-End pipeline from RAW data (DL0) to Science Products (DL4)
- FITS data format from DL0 to DL4
- C++/Python (pipeline and auxiliary modules)
- Single-telescope and array data reduction and analysis
- Additional components for data quality checks
- Monte Carlo data reduction and analysis (so far extensively tested)
- Real data reduction and analysis (so far tested with ASTRI-Horn telescope)
- Scientific analysis from DL3 to DL4 with external science tools (Gammapy and ctools)

# To appear soon

## Set of 4 papers to be published on the «Journal of High Energy Astrophysics»

### The ASTRI Mini-Array of Cherenkov Telescopes at the Observatorio del Teide

S. Scuderi<sup>a,\*</sup>, A. Giuliani<sup>a</sup>, G. Pareschi<sup>b</sup>, G. Tosti<sup>c</sup>, O. Catalano<sup>f</sup>, E. Amato<sup>p</sup>, L.A. Antonelli<sup>h</sup>, J. Becerra González<sup>m</sup>, G. Bellassai<sup>d</sup>, C. Bigongiari<sup>h</sup>, B. Biondo<sup>f</sup>, M. Boettcher<sup>n</sup>, G. Boranno<sup>d</sup>, P. Bruno<sup>d</sup>, A. Bulgarelli<sup>e</sup>, R. Canestrari<sup>f</sup>, M. Capalbi<sup>k</sup>, M. Cardillo<sup>k</sup>, V. Conforti<sup>e</sup>, C. Costantini<sup>f</sup>, M. Corpora<sup>f</sup>, A. Costa<sup>d</sup>, G. Cusumano<sup>f</sup>, A. D'Ai<sup>f</sup>, E. de Gouveia Dal Pino<sup>l</sup>, R. Della Ceca<sup>a</sup>, E. Escribano Rodríguez<sup>o</sup>, D. Falceta-Gonçalves<sup>s</sup>, C. Fermino<sup>l</sup>, M. Fiori<sup>h,g</sup>, G. Galanti<sup>o</sup>, G. Giacomini<sup>a</sup>, S. Gallozzi<sup>h</sup>, C. Gargano<sup>f</sup>, S. Garozzo<sup>d</sup>, S. Germani<sup>c</sup>, A. Ghedina<sup>o</sup>, F. Incardona<sup>q</sup>, M. J. Jarvis<sup>r</sup>, R. Jimenez<sup>f,l</sup>, V. Giordano<sup>d</sup>, A. Grillo<sup>d</sup>, C. Grivel Gelly<sup>o</sup>, D. Imr<sup>u</sup>, A. Incardona<sup>q</sup>, S. Incorvaia<sup>a</sup>, S. Iovenitti<sup>b</sup>, A. La Barbera<sup>f</sup>, N. La Palombara<sup>a</sup>, V. La Parola<sup>e</sup>, A. Lamastra<sup>h</sup>, L. Lessio<sup>g</sup>, G. Leto<sup>d</sup>, F. Lo Gerfo<sup>f</sup>, M. Lodi<sup>o</sup>, S. Lombardi<sup>h</sup>, F. Longo<sup>w,x</sup>, G. Lucarelli<sup>h</sup>, M.C. Maccarone<sup>f</sup>, D. Marano<sup>d</sup>, E. Martinetti<sup>d</sup>, S. Mereghetti<sup>a</sup>, A. Muccichè<sup>d</sup>, R. Millul<sup>b</sup>, T. Mineo<sup>f</sup>, G. Morlino<sup>o</sup>, A. Morselli<sup>i</sup>, G. Naletto<sup>j,g</sup>, G. Nicotra<sup>f</sup>, A. Pagliaro<sup>f</sup>, N. Parmiggiani<sup>e</sup>, G. Piano<sup>k</sup>, F. Pintore<sup>e</sup>, E. Poretti<sup>o</sup>, B. Olmi<sup>q</sup>, G. Rodeghiero<sup>e</sup>, G. Rodríguez Fernández<sup>i</sup>, P. Romano<sup>b</sup>, G. Romeo<sup>d</sup>, F. Russo<sup>e</sup>, P. Sangiorgi<sup>f</sup>, F.G. Saturni<sup>h</sup>, J.H. Schwarz<sup>b</sup>, E. Sciacca<sup>d</sup>, G. Sironi<sup>b</sup>, G. Sottile<sup>f</sup>, A. Stamerra<sup>h</sup>, G. Tagliaferri<sup>b</sup>, V. Testa<sup>h</sup>, G. Umana<sup>d</sup>, M. Uslenghi<sup>a</sup>, S. Vercellone<sup>b</sup>, L. Zampieri<sup>g</sup> and R. Zanmar Sanchez<sup>d</sup>

IN PRESS

### ASTRI Mini-Array Core Science at the *Observatorio del Teide*

S. Vercellone<sup>a,\*</sup>, C. Bigongiari<sup>b</sup>, A. Burtovoi<sup>c</sup>, M. Cardillo<sup>d</sup>, O. Catalano<sup>e</sup>, A. Franceschini<sup>f</sup>, S. Lombardi<sup>b,g</sup>, L. Nava<sup>a</sup>, F. Pintore<sup>e</sup>, A. Stamerra<sup>b</sup>, F. Tavecchio<sup>a</sup>, L. Zampieri<sup>h</sup>, R. Alves Batista<sup>i</sup>, E. Amato<sup>c,j</sup>, L. A. Antonelli<sup>b,g</sup>, C. Arcaro<sup>h,k</sup>, J. Becerra González<sup>l,m</sup>, G. Bonnoli<sup>a</sup>, M. Böttcher<sup>k</sup>, G. Brunetti<sup>n</sup>, A. A. Compagnino<sup>e</sup>, S. Cretan<sup>o,p</sup>, A. D'Ai<sup>e</sup>, M. Fiori<sup>h,f</sup>, G. Galanti<sup>o</sup>, A. Giuliani<sup>o</sup>, E. M. de Gouveia Dal Pino<sup>q</sup>, J. G. Green<sup>b</sup>, A. Lamastra<sup>b,g</sup>, M. Landoni<sup>a</sup>, F. Lucarelli<sup>b,g</sup>, G. Morlino<sup>c</sup>, B. Olmi<sup>r,c</sup>, E. Peretti<sup>s</sup>, G. Piano<sup>d</sup>, G. Ponti<sup>a,t</sup>, E. Poretti<sup>u</sup>, P. Romano<sup>a</sup>, F. G. Saturni<sup>h</sup>, S. Scuderi<sup>o</sup>, A. Tutone<sup>b</sup>, G. Umana<sup>v</sup>, J. A. Acosta-Pulido<sup>l,m</sup>, P. Barai<sup>q</sup>, A. Bonanno<sup>v</sup>, G. Boranno<sup>d</sup>, A. Bulgarelli<sup>w</sup>, V. Conforti<sup>w</sup>, A. Costa<sup>v</sup>, G. Cusumano<sup>e</sup>, M. Del Santo<sup>e</sup>, M. Della Ceca<sup>a</sup>, D. A. Falceta-Gonçalves<sup>q</sup>, V. Fioretti<sup>w</sup>, S. Germani<sup>x,y</sup>, R. J. Garcia-Cardena<sup>z</sup>, A. Ghedina<sup>u</sup>, V. Giordano<sup>v</sup>, M. Kreter<sup>k</sup>, F. Incardona<sup>v</sup>, S. Iovenitti<sup>a</sup>, A. La Barbera<sup>f</sup>, N. La Palombara<sup>o</sup>, V. La Parola<sup>e</sup>, G. Leto<sup>v</sup>, F. Longo<sup>z,aa</sup>, A. López-Oramas<sup>l,m</sup>, M. C. Maccarone<sup>f</sup>, S. Mereghetti<sup>o</sup>, R. Millul<sup>a</sup>, G. Naletto<sup>f</sup>, A. Pagliaro<sup>e</sup>, N. Parmiggiani<sup>w</sup>, C. Righi<sup>a</sup>, J. C. Rodríguez-Camírez<sup>q</sup>, G. Romeo<sup>v</sup>, P. Sangiorgi<sup>e</sup>, R. Santos de Lima<sup>q</sup>, G. Tagliaferri<sup>a</sup>, V. Testa<sup>b</sup>, G. Tosti<sup>x,y</sup>, M. Vázquez Acosta<sup>l,m</sup>, N. Żywucka<sup>k,ab</sup>, P. A. Caraveo<sup>o</sup> and G. Pareschi<sup>a</sup>

IN PRESS

### Galactic Observatory Science with the ASTRI Mini-Array at the *Observatorio del Teide*

A. D'Ai<sup>a,\*</sup>, E. Amato<sup>b</sup>, A. Burtovoi<sup>b</sup>, A. A. Compagnino<sup>a</sup>, M. Fiori<sup>c</sup>, A. Giuliani<sup>d</sup>, N. La Palombara<sup>d</sup>, A. Paizis<sup>d</sup>, G. Piano<sup>e</sup>, F. G. Saturni<sup>f,g</sup>, A. Tutone<sup>a,h</sup>, A. Belfiore<sup>d</sup>, M. C. Capalbi<sup>k</sup>, S. Cretan<sup>d</sup>, G. Cusumano<sup>a</sup>, M. Della Valle<sup>i,j</sup>, M. Del Santo<sup>a</sup>, A. La Barbera<sup>a</sup>, V. Conforti<sup>e</sup>, S. Lombardi<sup>f,g</sup>, S. Mereghetti<sup>d</sup>, G. Morlino<sup>b</sup>, F. Pintore<sup>a</sup>, P. Romano<sup>k</sup>, S. Vercellone<sup>b</sup>, L. Zampieri<sup>g</sup>, L. A. Antonelli<sup>h</sup>, C. Arcaro<sup>l</sup>, C. Bigongiari<sup>f</sup>, M. Böttcher<sup>m</sup>, P. Bruno<sup>n</sup>, A. Bulgarelli<sup>o</sup>, V. Conforti<sup>e</sup>, A. Costa<sup>n</sup>, E. de Gouveia Dal Pino<sup>p</sup>, V. Fioretti<sup>o</sup>, S. Germani<sup>q</sup>, A. Ghedina<sup>r</sup>, V. Giordano<sup>d</sup>, F. Incardona<sup>q</sup>, G. Leto<sup>n</sup>, F. Longo<sup>s,t</sup>, A. López Oramas<sup>u</sup>, F. Lucarelli<sup>f,g</sup>, B. Olmi<sup>v</sup>, A. Pagliaro<sup>f</sup>, N. Parmiggiani<sup>o</sup>, G. Romeo<sup>n</sup>, A. Stamerra<sup>f</sup>, V. Testa<sup>f</sup>, G. Tosti<sup>o,q</sup>, G. Umana<sup>n</sup>, L. Zampieri<sup>e</sup>, P. Caraveo<sup>d</sup> and G. Pareschi<sup>k</sup>

UNDER REVIEW

### Extragalactic Observatory Science with the ASTRI Mini-Array at the *Observatorio del Teide*

F. G. Saturni<sup>a,b,\*</sup>, C. H. E. Arcaro<sup>c,d,e,f</sup>, B. Balmaverde<sup>g</sup>, J. Becerra González<sup>h,i</sup>, A. Cacchione<sup>j</sup>, M. Capalbi<sup>k</sup>, A. Lamastra<sup>a</sup>, S. Lombardi<sup>a,b</sup>, F. Lucarelli<sup>a,b</sup>, R. Alves Batista<sup>l</sup>, L. A. Antonelli<sup>b</sup>, E. M. de Gouveia Dal Pino<sup>m</sup>, R. Della Ceca<sup>j</sup>, J. G. Green<sup>a,b</sup>, A. Pagliaro<sup>k</sup>, C. Righi<sup>a</sup>, F. Tavecchio<sup>n</sup>, S. Vercellone<sup>n</sup>, A. Wolter<sup>j</sup>, E. Amato<sup>o</sup>, C. Bigongiari<sup>a,b</sup>, M. Böttcher<sup>d</sup>, C. Costantini<sup>f</sup>, P. Bruno<sup>q</sup>, A. Bulgarelli<sup>r</sup>, M. Cardillo<sup>s</sup>, V. Conforti<sup>r</sup>, A. Costa<sup>q</sup>, G. Cusumano<sup>k</sup>, S. Germani<sup>t</sup>, A. Ghedina<sup>u</sup>, V. Giordano<sup>q</sup>, A. Giuliani<sup>v</sup>, F. Incardona<sup>q</sup>, A. La Barbera<sup>f</sup>, N. La Palombara<sup>o</sup>, G. Leto<sup>q</sup>, F. Longo<sup>w,x</sup>, G. Morlino<sup>o</sup>, B. Olmi<sup>y</sup>, N. Parmiggiani<sup>r</sup>, P. Romano<sup>n</sup>, G. Romeo<sup>v</sup>, A. Stamerra<sup>a</sup>, G. Tagliaferri<sup>n</sup>, V. Testa<sup>a</sup>, G. Tosti<sup>j,t</sup>, P. A. Caraveo<sup>v</sup> and G. Pareschi<sup>n</sup>

UNDER REVIEW

## ASTRI Mini-Array Core Science at the *Observatorio del Teide*

S. Vercellone<sup>a,\*</sup>, C. Bigongiari<sup>b</sup>, A. Burtovoi<sup>c</sup>, M. Cardillo<sup>d</sup>, O. Catalano<sup>e</sup>, A. Franceschini<sup>f</sup>, S. Lombardi<sup>b,g</sup>, L. Nava<sup>a</sup>, F. Pintore<sup>e</sup>, A. Stamerra<sup>b</sup>, F. Tavecchio<sup>a</sup>, L. Zampieri<sup>h</sup>, R. Alves Batista<sup>i</sup>, E. Amato<sup>c,j</sup>, L. A. Antonelli<sup>b,g</sup>, C. Arcaro<sup>h,k</sup>, J. Becerra González<sup>l,m</sup>, G. Bonnoli<sup>a</sup>, M. Böttcher<sup>k</sup>, G. Brunetti<sup>n</sup>, A. A. Compagnino<sup>e</sup>, S. Crestan<sup>o,p</sup>, A. D'Ai<sup>e</sup>, M. Fiori<sup>h,f</sup>, G. Galanti<sup>o</sup>, A. Giuliani<sup>o</sup>, E. M. de Gouveia Dal Pino<sup>q</sup>, J. G. Green<sup>b</sup>, A. Lamastra<sup>b,g</sup>, M. Landoni<sup>a</sup>, F. Lucarelli<sup>b,g</sup>, G. Morlino<sup>c</sup>, B. Olmi<sup>r,c</sup>, E. Peretti<sup>s</sup>, G. Piano<sup>d</sup>, G. Ponti<sup>a,t</sup>, E. Poretti<sup>a,u</sup>, P. Romano<sup>a</sup>, F. G. Saturni<sup>b,g</sup>, S. Scuderi<sup>o</sup>, A. Tutone<sup>b</sup>, G. Umana<sup>v</sup>, J. A. Acosta-Pulido<sup>l,m</sup>, P. Barai<sup>q</sup>, A. Bonanno<sup>v</sup>, G. Bonanno<sup>v</sup>, P. Bruno<sup>v</sup>, A. Bulgarelli<sup>w</sup>, V. Conforti<sup>w</sup>, A. Costa<sup>v</sup>, G. Cusumano<sup>e</sup>, M. Del Santo<sup>e</sup>, M. V. del Valle<sup>q</sup>, R. Della Ceca<sup>a</sup>, D. A. Falceta-Gonçalves<sup>q</sup>, V. Fioretti<sup>w</sup>, S. Germani<sup>x,y</sup>, R. J. García-López<sup>l,m</sup>, A. Ghedina<sup>u</sup>, V. Giordano<sup>v</sup>, M. Kreter<sup>k</sup>, F. Incardona<sup>v</sup>, S. Iovenitti<sup>a</sup>, A. La Barbera<sup>e</sup>, N. La Palombara<sup>o</sup>, V. La Parola<sup>e</sup>, G. Leto<sup>v</sup>, F. Longo<sup>z,aa</sup>, A. López-Oramas<sup>l,m</sup>, M. C. Maccarone<sup>e</sup>, S. Mereghetti<sup>o</sup>, R. Millul<sup>a</sup>, G. Naletto<sup>f</sup>, A. Pagliaro<sup>e</sup>, N. Parmiggiani<sup>w</sup>, C. Righi<sup>a</sup>, J. C. Rodríguez-Ramírez<sup>q</sup>, G. Romeo<sup>v</sup>, P. Sangiorgi<sup>e</sup>, R. Santos de Lima<sup>q</sup>, G. Tagliaferri<sup>a</sup>, V. Testa<sup>b</sup>, G. Tosti<sup>x,y</sup>, M. Vázquez Acosta<sup>l,m</sup>, N. Żywucka<sup>k,ab</sup>, P. A. Caraveo<sup>o</sup> and G. Pareschi<sup>a</sup>

<https://doi.org/10.1016/j.jheap.2022.05.005>

IN PRESS

The A  
del Te

S. Scude  
J. Becerr  
P. Bruno  
M. Corp  
E. Escrib  
S. Galloz  
R. Gime  
S. Incorv  
L. Lessic  
M.C. Ma  
G. Morli  
F. Pintor  
G. Rome  
G. Sottile  
L. Zamp

Galac  
Obse



A. D'Ai  
Palomba  
S. Crest  
S. Lomb  
C. Arcar  
Gouveia  
F. Long  
A. Stam

at the



W  
, E.  
chio<sup>n</sup>,  
mo<sup>q</sup>,  
ani<sup>t</sup>,  
go<sup>w,x</sup>,  
ferri<sup>n</sup>,

# The ASTRI Mini-Array – Performance

- We extend current IACTs **differential sensitivity up to several tens of TeV and beyond**
- Investigate possible spectral features at VHE, such as the presence of **spectral cut-offs** or the detection of emission at several tens of TeV expected from **Galactic PeVatrons**



# Angular and Energy resolution



# The ASTRI Mini-Array – Performance

PRELIMINARY	ASTRI Mini-Array	MAGIC	VERITAS	H.E.S.S.	HAWC	LHAASO	Tibet AS $\gamma$
Altitude [m]	2,390	2,396	1,268	1,800	4,100	4,410	4,300
FoV	$\sim 10^\circ$	$\sim 3.5^\circ$	$\sim 3.5^\circ$	$\sim 5^\circ$	2 sr	2 sr	2 sr
Angular Res.	0.05° (30 TeV)	0.07° (1 TeV)	0.07° (1 TeV)	0.06° (1 TeV)	0.15° (10 TeV)	(0.24–0.32)° (100 TeV)	$\sim 0.2^\circ$ (100 TeV)
Energy Res.	12% (10 TeV)	16% (1 TeV)	17% (1 TeV)	15% (1 TeV)	30% (10 TeV)	(13–36)% (100 TeV)	20% (100 TeV)
Energy Range	(0.3-200) TeV	(0.05-20) TeV	(0.08-30) TeV	(0.02-30) TeV	(0.1-200) TeV	(0.1-1,000) TeV	(0.1-1,000) TeV

## Sensitivity: better than current IACTs ( $E \gtrsim 3$ TeV)

Extended spectrum and cut-off constraints


## Energy/Angular resolution: $\sim 10\%$ / $\sim 0.05^\circ$ ( $E = 10$ TeV)

Characterize extended sources morphology


## $10^\circ$ field of view with homogeneous off-axis performance

Multi-target fields and extended sources


Enhanced chance for serendipitous discoveries



# Angular resolution and large field of view



ASTRI Mini-Array **200 hr simulation (up to  $E \sim 200$  TeV)** of the region **of the Galactic source 2HWC J1908+063**. The light green circle marks the  $\sim 0.52^\circ$  HAWC error-box for  $E > 56$  TeV



ASTRI Mini-Array **200 hr simulation of the Cygnus Region**. Green crosses mark the positions of the 3HWC sources in a  $10^\circ \times 10^\circ$  field of view

# The LHAASO PeVatrons

Cao et al., 2021, Nature

LHAASO Source	Possible Origin	Type	Distance (kpc)	Age (kyr) <sup>a</sup>	$L_s$ (erg/s) <sup>b</sup>	Potential TeV Counterpart <sup>c</sup>
LHAASO J0534+2202	PSR J0534+2200	PSR	2.0	1.26	$4.5 \times 10^{38}$	Crab, Crab Nebula
LHAASO J1825-1326	PSR J1826-1334	PSR	$3.1 \pm 0.2^d$	21.4	$2.8 \times 10^{36}$	HESS J1825-137, HESS J1826-130, 2HWC J1825-134
	PSR J1826-1256	PSR	1.6	14.4	$3.6 \times 10^{36}$	
LHAASO J1839-0545	PSR J1837-0604	PSR	4.8	33.8	$2.0 \times 10^{36}$	2HWC J1837-065, HESS J1837-069, HESS J1841-055
	PSR J1838-0537	PSR	$1.3^e$	4.9	$6.0 \times 10^{36}$	
LHAASO J1843-0338	SNR G28.6-0.1	SNR	$9.6 \pm 0.3^f$	$< 2^f$	—	HESS J1843-033, HESS J1844-030, 2HWC J1844-032
LHAASO J1849-0003	PSR J1849-0001	PSR	$7^g$	43.1	$9.8 \times 10^{36}$	HESS J1849-000, 2HWC J1849+001
	W43	YMC	$5.5^h$	—	—	
LHAASO J1908+0621	SNR G40.5-0.5	SNR	$3.4^i$	$\sim 10 - 20^j$	—	MGRO J1908+06, HESS J1908+063, ARGO J1907+0627, VER J1907+062, 2HWC 1908+063
	PSR 1907+0602	PSR	2.4	19.5	$2.8 \times 10^{36}$	
	PSR 1907+0631	PSR	3.4	11.3	$5.3 \times 10^{35}$	
LHAASO J1929+1745	PSR J1928+1746	PSR	4.6	82.6	$1.6 \times 10^{36}$	2HWC J1928+177, 2HWC J1930+188, HESS J1930+188, VER J1930+188
	PSR J1930+1852	PSR	6.2	2.9	$1.2 \times 10^{37}$	
	SNR G54.1+0.3	SNR	$6.3^{+0.8}_-0.7^d$	$1.8 - 3.3^k$	—	
LHAASO J1956+2845	PSR J1958+2846	PSR	2.0	21.7	$3.4 \times 10^{35}$	2HWC J1955+285
	SNR G66.0-0.0	SNR	$2.3 \pm 0.2^d$	—	—	
LHAASO J2018+3651	PSR J2021+3651	PSR	$1.8^{+1.7}_-1.4^l$	17.2	$3.4 \times 10^{36}$	MGRO J2019+37, VER J2019+368, VER J2016+371
	Sh 2-104	H II/YMC	$3.3 \pm 0.3^m / 4.0 \pm 0.5^n$	—	—	
LHAASO J2032+4102	Cygnus OB2	YMC	$1.40 \pm 0.08^o$	—	—	TeV J2032+4130, ARGO J2031+4157, MGRO J2031+41, 2HWC J2031+415, VER J2032+414
	PSR 2032+4127	PSR	$1.40 \pm 0.08^o$	201	$1.5 \times 10^{35}$	
	SNR G79.8+1.2	SNR candidate	—	—	—	
LHAASO J2108+5157	—	—	—	—	—	—
LHAASO J2226+6057	SNR G106.3+2.7	SNR	$0.8^p$	$\sim 10^p$	—	VER J2227+608, Boomerang Nebula
	PSR J2229+6114	PSR	$0.8^p$	$\sim 10^p$	$2.2 \times 10^{37}$	

Discovery of **12 sources emitting at several hundreds of TeV**, up to 1.4 PeV

Crab apart, the majority of remaining sources represent **diffuse  $\gamma$ -ray structures with angular extensions up to  $1^\circ$** , and all of them are located along the Galactic plane

The **actual sources** responsible for the ultra high-energy  $\gamma$ -rays **have not yet been firmly localized and identified** (except for the Crab Nebula), leaving open the origin of these extreme accelerators

The **ASTRI Mini-Array** will investigate these and future PeVatron sources, providing both the opportunity for **their precise identification** and important **information on their morphology**

# The Pillars' concept

**First 4 years**  
Specific science  
topics

**10° FoV**  
Several sources in a  
single pointing

**Pillar 1**  
The origin of cosmic rays

PeVatrons  
Particle propagation  
PWN HE emission  
UHECR from SB galaxies

**Pillar 2**  
Fundamental physics

IR EBL constraints  
Probing IGMF  
Blazars & hadron beams  
ALP & LIV

# Pillars' main scientific targets

## Pillar-1

Name	RA	Dec	Type	Zenith Angle <sup>1</sup>	Visibility <sup>2</sup>
PRELIMINARY	(deg)	(deg)		(deg)	(hr/yr)
Tycho	6.36	64.13	SNR	35.8	410+340
Galactic Center	266.40	-28.94	Diffuse	57.2	0+180
VER J1907+062	286.91	6.32	SNR+PWN	22	400+170
SNR G106.3+2.7	337.00	60.88	SNR	32.6	460+300
$\gamma$ -Cygni	305.02	40.76	SNR	12.5	460+160
W28/HESS J1800-240B	270.11	-24.04	SNR/MC	51.6	0+300
Crab	83.63	22.01	PWN	6.3	470+170
Geminga	98.48	17.77	PWN	10.5	460+170
M82	148.97	69.68	Starburst	41.4	310+470

## Pillar-2

Target	Class	RA (J2000)	DEC (J2000)	Obs. time	ZA	Moon	Strategy, analysis, notes
IAU Name				[hr]	[deg]	[%]	PRELIMINARY
IC 310	Radio gal.	03 16 43.0	+41 19 29	50-100	45	25	Better suited for ToO observations of high states
M87	Radio gal.	12 30 47.2	+12 23 51	50-100	45	25	Better suited for ToO observations of high states
Mkn 501	Blazar	16 53 52	+39 45 38	50-100	45	25	Better suited for ToO observations of high states

Target	Class	RA (J2000)	DEC (J2000)	Obs. time	ZA	Moon	Strategy, analysis, notes
IAU Name				[hr]	[deg]	[%]	
Mkn 501	Blazar	16 53 52.2	+39 45 36.6	50-100	45	25	LIV, ALP. Better suited for ToOs in high states.
1ES 0229+200	Blazar	02 32 48.6	+20 17 17.5	200	45	25	HB, LIV, ALP. Almost steady source, possible "fill in" target.
PRELIMINARY							

These lists of sources reflect the science knowledge **at the time of writing** this paper

We expect **to improve these lists** according to the new findings from both IACTs and PSAs

# The Galactic Center – a challenge in a challenge


It is a complex region harbouring several potential sources of particle acceleration

It can be observed by the ASTRI Mini-Array only at high zenith angles

Current IACTs detected **non-variable emission with no significant cut-off up to a few tens of TeV**

## ASTRI Mini-Array assets

- **the large FoV** will allow us to map the **whole GC region in a single observation**
- **the excellent angular resolution** could help us to **identify any HE source** among several candidates



Spatial and spectral characterization of the inner Galactic Ridge emission → (HESS Collab., 2018)

HESS, MAGIC and ASTRI spectra fitted with a proton population with a best fit cut-off at 120 PeV

**Exclude a cut-off in proton pop. below 3.5 PeV, 2.0 PeV, and 1.7 PeV at 68%, 90%, and 95% C.L.**

# Cosmic-ray propagation: $\gamma$ -Cygni


$\gamma$ -Cygni (G78.2+2.1) is a middle-aged SNR located in the Cygnus region and discovered by VERITAS

HAWC observed this source, but HAWC's low angular resolution does not allow one to drive firm conclusion on the spatial structure

We simulated **2 possible spectral models** (A and B) fitting the combined Fermi-LAT and VERITAS data

The ASTRI Mini-Array will **constrain** some physical parameters such as the **maximum energy reached by protons** and the **diffusion coefficient**

Moreover, it will **investigate the VHE emission morphology**



Black and red dots show the ASTRI Mini-Array simulations for model A and B, respectively, for 200 hr of exposure

# EBL studies in the IR regime

From the mid-IR to the far-IR, where the IR background intensity is maximal, EBL direct measurements are prevented by the overwhelming dominance of local emission from both the Galaxy and our Solar system

$$\lambda_{\max} \sim 1.24 \times E_{\text{TeV}} [\mu\text{m}]$$

Measurements in the **(10-30)TeV energy band probe the EBL in the  $\sim(10-30)\mu\text{m}$  regime**, otherwise inaccessible


**Best candidates to constrain the EBL up to  $\lambda \sim 100\mu\text{m}$ :**

**low-redshift radio galaxies**

M 87, IC 310, Centaurus A

**local star-bursting and active galaxies**

M 82, NGC 253, NGC 1068



Upper panel: extinction factor for photon-photon interaction on EBL at the IC 310 source distance.

Bottom panel: MAGIC (blue dots) and ASTRI Mini-Array (red dots) 50 hours,  $5\sigma$  differential sensitivity


# Fundamental physics – hadron beams

Relativistic jets from extreme BL Lacs could be one of the UHECR acceleration sites

**Jets in extreme BL Lac objects could produce hadron beam (collimated beams of high-energy protons/nuclei)**

While travelling towards the Earth

- UHECR lose energy through photo-meson and pair production
- these trigger the development of electromagnetic cascades producing  $\gamma$  and  $\nu$ .
- Because of the reduced distance,  $\gamma$  experience a less severe EBL absorption
- **The observed gamma-ray spectrum extends at energies much higher ( $E > 10\text{TeV}$ ) than those allowed by the conventional EBL propagation**



Simulated VHE spectrum of 1ES 0229+220 for the standard (light blue, 200 hr) and hadron beam (red, 250 hr) scenarios

**The ASTRI Mini- Array would be able to obtain a significant detection up to 20 TeV with a deep (~250 hr) observation**


# Gamma-ray bursts

- GRBs confirmed as a new class of TeV emitters thanks to the MAGIC detection of GRB 190114C ( $z=0.42$ )
- SSC component extending into the TeV energy range

## The ASTRI Mini-Array

- might have detected emission from GRB 190114C
- is able to confirm afterglow emission at  $E > 1$  TeV from close ( $z < 0.4$ ) GRBs if observations start within the first tens of seconds up to few minutes from the onset of the burst
- can measure the spectral cut-off, either originated by the EBL absorption or intrinsic, if greater than 1 TeV

**The expected number of follow-ups on observable GRBs is about than 1 per month**



Simulation of the emission from three GRB 190114C-like bursts, at three different redshifts ( $z = 0.078$ ,  $z = 0.25$  and  $z = 0.42$ )

# Strategic VHE synergies

- Both **MAGIC** and **CTAO-N** will be of paramount importance for the **study of GRBs**, as will be their capability to investigate not only the local Universe, but also reaching **redshifts well beyond one**
- Both **MAGIC** and **CTAO-N** will allow us to extend the ASTRI Mini-Array spectral performance in the **sub-TeV regime**, with almost no breaks **from a few tens of GeV up to hundreds of TeV**
- The **PSAs** detected several sources with **photons up to several hundreds of TeV**. Potential synergies are important to make use of the **ASTRI Mini-Array angular and energy resolution** in combination with the LHAASO, HAWC and Tibet AS $\gamma$  extended energy range

# The multi-wavelength landscape

- **MeerKat** and **ASCAP** (SKA precursors in the South) will allow us to investigate the Galactic Center and its features
- **LOFAR** (SKA precursor in the North) will open a new science window in the low-frequency radio band and monitor 2/3 of the sky nightly in Radio Sky Monitor mode, being an excellent radio transient factory
- **SRT** has already observed sources of interest for the ASTRI Mini-Array, such as W 44, IC 433 and Tycho, making it an excellent observatory for future synergies in the northern hemisphere
- **TNG** is located in La Palma and can be extremely useful for optical follow-up observations. The **WEBC Consortium** is dedicated to the observation of blazars, and it is fundamental for blazar SEDs. IAC also provides access to several optical telescopes on-site.
- **eROSITA/SRG, XMM-Newton, Chandra, NuSTAR and IXPE** will provide fundamental photometric, imaging, spectroscopic, and polarimetric data.
- **AGILE, Fermi, INTEGRAL, and Swift** will be extremely important for their large FoV and for the *Swift* ability to promptly react to transients

# Core Science and Observing Plan


## Baseline

- About 1500 moonless hours/year at the Teide site
- Bad weather, maintenance, calibrations... → **~1000 hours/year for scientific observations**

## Room for improvement

- ASTRI Mini-Array camera composed of SiPM, → **observations with a significant fraction of the Moon**, in addition to the 1000 hr/yr
- Main scientific goals focus on the multi-TeV energy band → we can **effectively perform observations at high (~60°) zenith angles**

Sources	Season	Dark hours
Galactic Center	May – June – July	300
VER J1907+062	September – October	300
G106.3+2.7	November – December	400



This example shows that we can observe **several sources per year** thanks to their different sky positions

We expect also **serendipitously detected sources**, thanks to the ASTRI Mini-Array wide field of view

# From Core Science to Observatory Science

---

For the first 4 years the ASTRI Mini-Array will be run as an experiment

It will be dedicated to the Core Science Topics

**Smooth transition towards an Observatory period**

**Build-up on the experience and results from the Core Science**

**Open to observational proposals from the scientific community**

The ASTRI Mini-Array will start **scientific observations in 2024** from the *Observatorio del Teide* with a 4 (core science) + 4 (observatory science) year programme

Its **10° field of view** will allow us to investigate both extended sources (e.g., SNRs) and crowded/rich fields (e.g., the Galactic Center) with a single pointing

Its **3' angular resolution** at 10 TeV will allow us to perform detailed morphological studies of extended sources

Its **sensitivity extending above 100 TeV** will make it the most sensitive IACT in the energy range 5-200 TeV in the Northern hemisphere

It will **join together** the **very high-energy domain** typical of PSAs with the **precision domain** (excellent angular and energy resolutions) typical of IACTs