


Multiphase AGN-driven outflows and X-ray winds

Elena Bertola
X-ray Lab AA 2021-2022

Why do we care about AGN feedback?




1. Galaxy evolution models


The mass functions of **dark matter haloes** and **galaxies** don't agree at the *low and high mass ends*

Part of the gas available in the galaxy is not converted in stars

Why?



Why do we care about AGN feedback?



1. Galaxy evolution models

Models can explain the low-mass end if Supernova feedback is accounted for

Similarly, the high-mass end can be matched including AGN feedback


Why do we care about AGN feedback?

2. SMBH/host-galaxy scaling relations


Are AGN and host galaxies coevolving?

How is this coevolution set into place?

Magorrian+1989, Gebhardt+2000, Ferrarese+2000,2006, McConnell+2011, Kormendy & Ho 2013, etc.



Alexander & Hickox, 2018



A logarithmic view of the AGN-galaxy connection


Binding Energies

$$E_{b,\odot} \approx 4 \times 10^{48} \text{ ergs}$$


$$E_{b,BH,8} \approx 10^{61} \text{ ergs}$$

$$E_{b,gal,11} \approx 10^{59} \text{ ergs}$$

$$E_{b,Coma} \approx 10^{64} \text{ ergs}$$




Types of AGN feedback



Radio mode: AGN relativistic jets


- Accretion onto the BH is generally inefficient
- Little energy goes in radiation
- Great part of the energy produced by the AGN is converted in kinetic energy
→ *relativistic jets*
- Predominantly found in the most massive galaxies ($M_{\text{star}} > 10^{11} M_{\odot}$) with old stellar populations
- Jets heat up the gas they encounter, creating bubbles and cavities and *preventing radiative cooling*
- Radio mode can explain why we don't find cooling flows in galaxy clusters




< ————— 700 kpc ————— >

Quasar mode: AGN-driven winds

- Accretion onto the BH is efficient, Eddington ratios are high
- Great part of the AGN energy goes in radiation
- Radiative AGN are most common in galaxies with on-going star-formation and younger stellar populations at all cosmic epochs
- Production of AGN-driven *winds* via
 - Radiation driving
 - Line driving
 - Magnetic acceleration
- Impact on the SF activity by
 - heating
 - dissociating
 - removingthe cold gas reservoir of the host galaxy



Modeling feedback through AGN-driven winds



$$\dot{E}_{\text{kin}}^{\text{wind}} / L_{\text{bol}} = 0.05$$


Di Matteo+2005:

The observed scaling relations can be matched by simulations if the energy-loading factor of the AGN outflows is at least 5%

e.g., King 2003, Di Matteo+2005, Hopkins+2006, Hopkins&Elvis2010, Faucher-Giguère&Quataert 2012, Lapi+2014, Costa+2018,2020, ...

How are AGN-driven winds linked to kpc-scale outflows?

Inner-disc wind is launched with velocity v_{in}




Momentum-driven
vs.
Energy-driven

Inner-disc wind propagation


Inner-disc wind shocks the ISM *only if* the mechanical luminosity of the wind is thermalized at R_c , i.e., cooling at R_c is negligible

How are AGN-driven winds linked to kpc-scale outflows?



From the observational side, we want to populate this diagram and check if the winds we observe match any of the two expectations

How are AGN-driven winds linked to kpc-scale outflows?




From the observational side, we want to populate this diagram and check if the winds we observe match any of the two expectations

Inner-disc winds

Inner-disk winds: Ultra-fast Outflows (UFOs)

The *fastest* winds known to date



Present in 40% of RQ *and* RL AGN

(Tombesi+10,+14,Gofford+13,Igo+20)

$N_H \sim 10^{22}-10^{24}$ cm^{-2} \rightarrow thick

$\log(\xi/erg\ s^{-1}cm^2) > 3$ \rightarrow highly ionized

$v > 0.05c$ (up to $0.6c$) \rightarrow ultra fast

Signature: P-cygni profile of FeXXV-XXVI

Commonly seen as resonant absorption lines of highly-ionized iron blueshifted at $E > 7$ keV

Need high S/N spectra to constrain them

Observing AGN-driven winds

Inner to outer:

1. UFOs, WA
2. UV BAL–NAL

} Disc winds

Narrow Absorption Lines (NAL)
 $\log \xi = 0 - 1.5 \text{ erg cm s}^{-1}$
 $\log N_H = 18 - 20 \text{ cm}^{-2}$
 Velocity = 100 - 1000 km/s
 Distance scale $\sim 1 \text{ pc} - 1 \text{ kpc}$



Broad Absorption Line (BAL)
 $\log \xi = -0.5 \text{ to } 2.5 \text{ erg cm s}^{-1}$
 $\log N_H = 20 - 23 \text{ cm}^{-2}$
 Velocity = 10,000- 60,000 km/s
 Distance scale = 0.001 pc - 500 pc

X-rays

Warm Absorbers (WA)
 $\log \xi = -1 \text{ to } 3.0 \text{ erg cm s}^{-1}$
 $\log N_H = 21 \text{ to } 22.5 \text{ cm}^{-2}$
 Velocity = 100 - 2000 km/s
 Distance scale = 0.1 pc - 1 kpc

Ultra Fast Outflow (UFO)
 $\log \xi = 3 - 5.0 \text{ erg cm s}^{-1}$
 $\log N_H = 22 - 23.5 \text{ cm}^{-2}$
 Velocity = 10,000- 70,000 km/s
 Distance scale = 0.001 pc - 10 pc

UV



Laha+2020 (Review)

Observing AGN-driven winds

Inner to outer:

1. UFOs, WA
 2. UV BAL–NAL
- } Disc winds

3. Ionized outflows (e.g., [OIII])
4. Neutral gas outflows
5. Molecular outflows

Galaxy-wide outflows




Observing AGN-driven winds


Inner to outer:

1. UFOs, WA
 2. UV BAL–NAL
- } Disc winds

3. Ionized outflows (e.g., [OIII])
4. Neutral gas outflows
5. Molecular outflows

Galaxy-wide outflows






W_{80} : a measure of the outflow strength

Non parametric approach

v_n :
n-th percentile of the
emission-line profile
i.e., n% of the line area is
enclosed at the left of v_n

$$W_{80} = v_{90} - v_{10}$$




Observing AGN-driven winds

Inner to outer:

1. UFOs, WA
 2. UV BAL–NAL
- } Disc winds

3. Ionized outflows (e.g., [OIII])
4. Neutral gas outflows
5. Molecular outflows

Galaxy-wide outflows




Observing AGN-driven winds

Inner to outer:

1. UFOs, WA
 2. UV BAL–NAL
- } Disc winds

3. Ionized outflows (e.g., [OIII])
4. Neutral gas outflows
5. Molecular outflows

Galaxy-wide outflows



Local and low- z AGN

Testing the models with observations — I

Issue:


Need for **multi-wavelength coverage** and detection of multiphase outflows

Only a handful of AGN were found to show inner-disc- *and* galaxy-scale outflows

Workaround:

Build big samples of AGN outflows and search for correlations *within* the sample


e.g.: Fiore+2017 – the outflow velocity correlates with the AGN L_{bol} for **molecular+ionized** outflows and **UFOs**, and that the two scalings are statistically consistent with each other



Mout vs. Lbol is flatter for molecular outflows

Most powerful AGN seem to have more massive outflows in the ionized phase

Need to measure all the gas phases, especially at high luminosities, to properly study the impact of AGN winds




(a)

Energy-conserving scenario:




- - Bulk of the outflow mass is molecular

□ Luminosity corrected - a significant part of the outflow mass is in the ionized phase



We can test the impact of AGN outflows by **comparing the mass-outflow rate to the SFR**, that is comparing how much gas is removed by the outflow to how much gas is converted into stars


mass-loading factor > 1: the outflow removes the gas faster than it takes to form stars – negative AGN feedback



AGN winds are more effective in suppressing the SF than winds driven by starburst activity

Mass-loading factor shows mild correlation with AGN luminosity or SMBH mass

Testing the models with observations — II




Another interesting parameter is the energy loading factor, because we can directly compare our results with the model predictions – *but can we??*

Issue 1 :

we need some assumptions to compute the kinetic power of the winds (e.g., geometry of the wind, gas density)

Testing the models with observations — II




Issue 2:
models do not agree on the predicted threshold for the coupling efficiency

Issue 3:
model predictions are usually based on all the outflowing material and not on the individual gas phases — back to the argument of Bischetti+19


AGN feedback at high- z

High-redshift AGN and AGN outflows




Local and low- z AGN allow for detailed studies of spatially resolved outflows and gas dynamics

But is the local Universe where we expect AGN feedback to be at its highest?



High-redshift AGN and AGN outflows



Star formation history

Black Hole Accretion history:

Shankar+2009

Aird+2010




Del Vecchio+2014

Here is where we expect AGN feedback to be at its maximum, but observations are more challenging than in the local Universe

High-z UFOs

Local samples:
tens of AGN

$V_{\text{mean}} \sim 0.15$
 $DF \sim 0.3-0.4$





Chartas+2003,+2007,+2009b,+2016+2021, Vignali+2015,
Dadina+2018, Lanzuisi+2012, Bertola+2020

Chartas+21: XMM program on NAL AGN ■


NAL AGN show a UFO DF~80%
hints at link between multiphase winds on the
meso- and micro-scale

However, there is no clear link between the
velocities of the two components



Chartas+2003,+2007,+2009b,+2016+2021, Vignali+2015,
Dadina+2018, Lanzuisi+2012, Bertola+2020



The case of XID2028 ($z=1.593$): positive and negative feedback, and CO depletion



Narrow $H\alpha$ map:
tracer of SF

Black contours:
[OIII] outflow


Cresci+2015



Brusa+2018

Black contours:
molecular outflow

Green contours:
[OIII] outflow



obscured QSOs:

- Aravena et al. (2008)
- Feruglio et al. (2014)
- Popping et al. (2017)
- * Mrk 231 (Fiore et al. 2017)

reference samples:

- $0 < z < 2.5$ MS galaxies (Sa14)
- × $z \sim 0$ starburst (U)LIRGs (So97)
- $z \sim 1.5$ off-MS starbursts (Si15)
- $1.2 < z < 3.4$ SMGs (Bo13)
- $1 < z < 1.5$ AGN (Kakkad et al. 2017)

literature scaling relations

($z=1.6$ & $10.7 < \log(M_*/M_\odot) < 11.6$):

- Tacconi et al. (2017)
- Scoville et al. (2017)
- Sargent et al. (2014)

...but does AGN feedback actually have an impact on the star formation of galaxies?

The impact of AGN feedback on the SF of galaxies

If AGN winds establish the SMBH/host coevolution, then **AGN must act on the SF of the host galaxy.**

The most straightforward way to **act on** the SF is to impact the **fuel of (future) SF**, that is the molecular gas content

AGN are thus expected to reduce the molecular gas content of galaxies, but **do we see this in our data?**
And what do the simulations predict?



The impact of AGN feedback on the SF of galaxies

How do we study this:





1. Build **samples of AGN**, measure the **total molecular gas content** of their host galaxies and obtain the **other properties of the galaxy from SED fitting** (for instance, the stellar mass, SFR).
2. Search for **correlations** between the **total molecular gas mass** of the host galaxy and molecular and/or ionized **outflows** (if any) and with AGN properties
3. **Build a control sample of non-active galaxies** matched in SFR and/or stellar mass to the AGN sample and search for differences - e.g., are AGN CO depleted?

CO depletion: at fixed SFR/stellar mass, one galaxy sample shows *less* molecular gas than the selected control sample

Low redshift and local Universe

AGN host galaxies are consistent with non-active galaxies


The more powerful the AGN, the *more gas rich* the host:
AGN need gas to power the central activity,
the same gas that powers star formation



High redshift

High-z AGN seem to reside in CO depleted host galaxies

Issue: studies based on few, powerful AGN, pre-selected as good candidates for hosting outflows




KASHz and SUPER

SUPER: *SINFONI* Survey for Unveiling the Physics and Effect of Radiative feedback


Unbiased samples of high- z , X-ray selected AGN spanning a wide range of AGN power

Aim: study relation between AGN-driven ionized outflows and host galaxy properties (star formation and molecular gas content)

KASHz: *KMOS* AGN Survey at High redshift



KASHz and SUPER outflows



KASHz AGN:

strong [OIII] outflows + targets
consistent to non-active galaxies

SUPER AGN Type 1:



all have $w_{80} > 600$ km/s

Samples are matched in
redshift and stellar mass


Kakkad+2020

Difference between
KASHz and SUPER:
X-ray luminosity

SUPER OFs match
 $\dot{M}_{\text{out}} - L_{\text{bol}}$ relations of
local and low-z AGN




SUPER AGN - CO depletion?




Demonstrates the importance of including 'regular' AGN in our high-z samples

SUPER AGN show significant CO depletion only in the most massive host galaxies ($M_{\text{star}} > 10^{11} M_{\odot}$)



And what do simulations predict?




Ward+2022,
subm.

AGN live in high gas fraction
and high SFR galaxies

Qualitative agreement with results in
low-z AGN

Potentially in tension with those at
high-z, which also have a less
coherent picture

And what do simulations predict?



Ward+2022, subm.

Predictions on gas-depletion are **simulation-dependent**, but in general there is **no evidence for AGN hosts being more depleted** than non-active galaxies

Issues:

1. It is hard for cosmological simulations to implement small-scale AGN feedback
→ need for smaller grids and better physical implementation of feedback
2. Include results on high- L_{bol} AGN (rare and short lived)
→ need for bigger volumes



CIAO

CIAO



Piace a [alessandropeca](#) e altre persone

[astromemes_unibo](#) E dal festival di Quasan Remo è tutto 🎵