


XMM-Newton data analysis: tutorial





Eleonora Torresi
(INAF-OAS Bologna)


Outline

- The spacecraft
- Webpage, software and data download
- Data structure
- SAS
- Data re-processing
- Data reduction:
 - filtering for high particle background
 - selection of good time intervals (GTI)
 - selection of source and background extraction regions
- EPIC science modes
- Pileup
- Extraction of source and background spectra
- RMF & ARF
- Grouping
- (Background corrected) light curve extraction

The spacecraft



Highly elliptical orbit around the Earth
Inclination 40 degrees to the Equator



See M. Dadina's presentation & C. Vignali's presentation

1. XMM-Newton archive & data download

<https://www.cosmos.esa.int/web/xmm-newton/xsa>

XMM-Newton » Archive, Pipeline & Catalogues » XMM-Newton Science Archive

Home / Latest News
XMM-Newton 20th Anniversary
Conferences & Meetings
News
General User Support
Proposers Info
Observers Info
Data Analysis
Archive, Pipeline & Catalogues
Calibration & Background
SOC Info
About XMM-Newton
Image Gallery
Publications
Other Links

XMM-NEWTON SCIENCEX ARCHIVE (XSA)

INDEX

- Access to XMM-Newton Data and Source Catalogues
- Download Full XMM-Newton Catalogues and datasets **New**
- Tools
- Watchouts
- Notes on the XSA releases **New**
- Documentation
- Questions, Comments

ACCESS TO XMM-NEWTON DATA AND SOURCE CATALOGUES


[Search the XMM-Newton Science Archive \(XSA\)](#)

Direct access to the XSA data via URL or AIO (Archive InterOperability System):

[Command line and URL access to the XSA data](#)

Astroquery and TAP (Table Access Protocol) access to the XSA Database:

[Astroquery and TAP queries to the XSA Database](#)





XMM-Newton Science Archive Search

Position File

- Name
- Equatorial
- Galactic

Target in Field Of View Circle Box

Name

3C 33

for Resolve

► Observation and Proposal filters

► Display options

[Reset Form](#)

[Catalogue Search >](#)

 [Submit](#)

XMM-Newton Science Archive



HOME SEARCH COMMAND & URL ACCESS INTERACTIVE ANALYSIS TAP QUERIES ASTROQUERY

Basket

Back to Search Close all

Results #1

OBSERVATIONS (2)

			Obs.ID	EPIC	RGS	BKGD	ESASky	Target	RA	DEC	Rev	Distance	Start Date	End Date	Dur.	Target Type	PI name	Prop. Program	P
<input type="checkbox"/>			0203280101					3C33	01h 08m 52.85s	+13d 20' 13.8"	745	0.01	2004-01-04 04:49:31	2004-01-04 08:59:23	14992	RADIO GALAXY RADIO LOUD/STEEP SPECT	HARDCASTLE, MARTIN	GO	I
<input checked="" type="checkbox"/>			0203280101					3C33	01h 08m 52.85s	+13d 20' 13.8"	754	0.01	2004-01-21 17:06:11	2004-01-21 19:34:37	8906	RADIO GALAXY RADIO LOUD/STEEP SPECT	HARDCASTLE, MARTIN	GO	I

- [ODF](#)
- [PPS](#)
- [IMAGES](#)
- [SOURCES](#)
- [SPECTRA](#)
- [LIGHT_CURVES](#)

Public Date	PPS ver	Coord. Obs
Public data	17.56_20190403_1200	-
Public data	17.56_20190403_1200	-

XMM-Newton Science Archive



HOME SEARCH COMMAND & URL ACCESS INTERACTIVE ANALYSIS TAP QUERIES ASTROQUERY


Basket

Back to Search Close all

Results #1

OBSERVATIONS (2)

<input type="checkbox"/>			Obs.ID	EPIC	RGS	BKGD	ESASky	Target	RA	DEC	Rev	Distance	Start Date	End Date
<input type="checkbox"/>			0203280101					3C33	01h 08m 52.85s	+13d 20' 13.8"	745	0.01	2004-01-04 04:49:31	2004-01-04 08:59:23
<input checked="" type="checkbox"/>			0203280301					3C33	01h 08m 52.85s	+13d 20' 13.8"	754	0.01	2004-01-21 17:06:11	2004-01-21 19:34:37



Summary Science Exposures Publications

Obs. ID	0203280301
Revolution	754
Target	3C33
Exposures	3 EPIC, 1 OM, 2 RGS

Proposal Abstract

A SEARCH FOR THE GROUP ENVIRONMENTS OF LOW-POWER FRII RADIO GALAXIES

Although a good deal of indirect evidence points towards moderate or poor groups as the environment of low-luminosity (and therefore typical) FRII radio galaxies, there are almost no direct X-ray observations of the required hot, confining medium. An FRII radio source would have a dramatic effect on the gas in a group, heating and potentially expelling a significant fraction of it, with important consequences for our understanding of group evolution. In order to characterize the X-ray environments of typical FRII sources, we propose a small survey of the nearest few FRII sources of intermediate size drawn from the well-studied 3CRR catalogue. Our observations will allow us to detect, and measure the properties of, group-scale emission around our targets.

[Show Quality Report](#)

XMM-Newton Science Archive



HOME SEARCH COMMAND & URL ACCESS INTERACTIVE ANALYSIS TAP QUERIES ASTROQUERY

Basket

Back to Search Close all X

Results #1 X

OBSERVATIONS (2) X

			Obs.ID	EPIC	RGS	BKGD	ESASky	Target	RA	DEC	Rev	Distance	Start Date	End Date	Dur.	Target Type	PI name	Prop. Program	P
<input type="checkbox"/>			0203280101					3C33	01h 08m 52.85s	+13d 20' 13.8"	745	0.01	2004-01-04 04:49:31	2004-01-04 08:59:23	14992	RADIO GALAXY RADIO LOUD/STEEP SPECT	HARDCASTLE, MARTIN	GO	I
<input checked="" type="checkbox"/>			0203280101					3C33	01h 08m 52.85s	+13d 20' 13.8"	754	0.01	2004-01-21 17:06:11	2004-01-21 19:34:37	8906	RADIO GALAXY RADIO LOUD/STEEP SPECT	HARDCASTLE, MARTIN	GO	I

- ODF
- PPS
- IMAGES
- SOURCES
- SPECTRA
- LIGHT_CURVES

odf= observation data files => raw data

0203280301.tar.gz

Data structure

0203280301.tar.gz → tar -zxvf 0203280301.tar.gz

**-z= unzip
-xvf=untar**

0231_0065940101.TAR	0231_0065940101_0MS40100RFX.FIT	0231_0065940101_0MU00700WDX.FIT	0231_0065940101_PNU00706DLI.FIT	0231_0065940101_R1S91904DII.FIT	0231_0065940101_R2S91509DII.FIT
0231_0065940101_M1S00300AUX.FIT	0231_0065940101_0MS40100THX.FIT	0231_0065940101_0MU00701IMI.FIT	0231_0065940101_PNU00807DLI.FIT	0231_0065940101_R1S92005DII.FIT	0231_0065940101_R2S91601DII.FIT
0231_0065940101_M1S00310IME.FIT	0231_0065940101_0MS40100WDX.FIT	0231_0065940101_0MU00800IMI.FIT	0231_0065940101_PNU00908DLI.FIT	0231_0065940101_R1S92106DII.FIT	0231_0065940101_R2S91702DII.FIT
0231_0065940101_M1S00320IME.FIT	0231_0065940101_0MS40101IMI.FIT	0231_0065940101_0MU00800WDX.FIT	0231_0065940101_PNU01009DLI.FIT	0231_0065940101_R1S92208DII.FIT	0231_0065940101_R2S91803DII.FIT
0231_0065940101_M1S00330IME.FIT	0231_0065940101_0MS40200IMI.FIT	0231_0065940101_0MU00801IMI.FIT	0231_0065940101_PNU01110DLI.FIT	0231_0065940101_R1S92309DII.FIT	0231_0065940101_R2S91905DII.FIT
0231_0065940101_M1S00340IME.FIT	0231_0065940101_0MS40200RFX.FIT	0231_0065940101_0MX00000NPH.FIT	0231_0065940101_PNU01211DLI.FIT	0231_0065940101_R1S92401DII.FIT	0231_0065940101_R2S92006DII.FIT
0231_0065940101_M1S00350IME.FIT	0231_0065940101_0MS40200THX.FIT	0231_0065940101_0MX00000PEH.FIT	0231_0065940101_PNU01312DLI.FIT	0231_0065940101_R1S92502DII.FIT	0231_0065940101_R2S92107DII.FIT
0231_0065940101_M1S00360IME.FIT	0231_0065940101_0MS40200WDX.FIT	0231_0065940101_PNS00400AUX.FIT	0231_0065940101_PNX00000HCH.FIT	0231_0065940101_R1S92603DII.FIT	0231_0065940101_R2S92208DII.FIT
0231_0065940101_M1S00370IME.FIT	0231_0065940101_0MS40201IMI.FIT	0231_0065940101_PNS00400CCX.FIT	0231_0065940101_PNX00000PAH.FIT	0231_0065940101_R1S92704DII.FIT	0231_0065940101_R2S92309DII.FIT
0231_0065940101_M1X00000HBH.FIT	0231_0065940101_0MS40300IMI.FIT	0231_0065940101_PNS00401IME.FIT	0231_0065940101_PNX00000PMH.FIT	0231_0065940101_R1S92805DII.FIT	0231_0065940101_R2S92401DII.FIT
0231_0065940101_M1X00000HCH.FIT	0231_0065940101_0MS40300RFX.FIT	0231_0065940101_PNS00402IME.FIT	0231_0065940101_R1S00500AUX.FIT	0231_0065940101_R1X00000D1H.FIT	0231_0065940101_R2S92502DII.FIT
0231_0065940101_M1X00000HTH.FIT	0231_0065940101_0MS40300THX.FIT	0231_0065940101_PNS00403IME.FIT	0231_0065940101_R1S00501SPE.FIT	0231_0065940101_R1X00000D2H.FIT	0231_0065940101_R2S92603DII.FIT
0231_0065940101_M1X00000PEH.FIT	0231_0065940101_0MS40300WDX.FIT	0231_0065940101_PNS00404IME.FIT	0231_0065940101_R1S00502SPE.FIT	0231_0065940101_R1X00000FX.FIT	0231_0065940101_R2S92705DII.FIT
0231_0065940101_M1X00000PTH.FIT	0231_0065940101_0MS40301IMI.FIT	0231_0065940101_PNS00405IME.FIT	0231_0065940101_R1S00503SPE.FIT	0231_0065940101_R1X00000PFH.FIT	0231_0065940101_R2S92806DII.FIT
0231_0065940101_M2S00700AUX.FIT	0231_0065940101_0MS40400IMI.FIT	0231_0065940101_PNS00406IME.FIT	0231_0065940101_R1S00504SPE.FIT	0231_0065940101_R2S00600AUX.FIT	0231_0065940101_R2X00000D1H.FIT
0231_0065940101_M2S00710IME.FIT	0231_0065940101_0MS40400RFX.FIT	0231_0065940101_PNS00407IME.FIT	0231_0065940101_R1S00505SPE.FIT	0231_0065940101_R2S00601SPE.FIT	0231_0065940101_R2X00000D2H.FIT
0231_0065940101_M2S00720IME.FIT	0231_0065940101_0MS40400THX.FIT	0231_0065940101_PNS00408IME.FIT	0231_0065940101_R1S00506SPE.FIT	0231_0065940101_R2S00602SPE.FIT	0231_0065940101_R2X000000FX.FIT
0231_0065940101_M2S00730IME.FIT	0231_0065940101_0MS40400WDX.FIT	0231_0065940101_PNS00409IME.FIT	0231_0065940101_R1S00508SPE.FIT	0231_0065940101_R2S00603SPE.FIT	0231_0065940101_R2X00000PFH.FIT
0231_0065940101_M2S00740IME.FIT	0231_0065940101_0MS40401IMI.FIT	0231_0065940101_PNS00410IME	0231_0065940101_R1S00509SPE.FIT	0231_0065940101_R2S00605SPE.FIT	0231_0065940101_RMX00100ECX.FIT
0231_0065940101_M2S00750IME.FIT	0231_0065940101_0MS40500IMI.FIT	0231_0065940101_PNS00411IME	0231_0065940101_R1S90001DII.FIT	0231_0065940101_R2S00606SPE.FIT	0231_0065940101_RMX00100ESX.FIT
0231_0065940101_M2S00760IME.FIT	0231_0065940101_0MS40500RFX.FIT	0231_0065940101_PNS00412IME	0231_0065940101_R1S90102DII.FIT	0231_0065940101_R2S00607SPE.FIT	0231_0065940101_SCX00000ATS.FIT
0231_0065940101_M2S00770IME.FIT	0231_0065940101_0MS40501IMI.FIT	0231_0065940101_PNU00200IMI.FIT	0231_0065940101_R1S90203DII.FIT	0231_0065940101_R2S00608SPE.FIT	0231_0065940101_SCX00000P1S.FIT
0231_0065940101_M2X00000HBH.FIT	0231_0065940101_0MS40400WDX.FIT	0231_0065940101_PNU00200IMI.FIT	0231_0065940101_R1S90304DII.FIT	0231_0065940101_R2S00609SPE.FIT	0231_0065940101_SCX00000P2S.FIT
0231_0065940101_M2X00000HCH.FIT	0231_0065940101_0MS40400THX.FIT	0231_0065940101_PNU00200WDX.FIT	0231_0065940101_R1S90405DII.FIT	0231_0065940101_R2S90001DII.FIT	0231_0065940101_SCX00000P3S.FIT
0231_0065940101_M2X00000HTH.FIT	0231_0065940101_0MS40400WDX.FIT	0231_0065940101_PNU00201IMI.FIT	0231_0065940101_R1S90506DII.FIT	0231_0065940101_R2S90102DII.FIT	0231_0065940101_SCX00000P4S.FIT
0231_0065940101_M2X00000PEH.FIT	0231_0065940101_0MS40400THX.FIT	0231_0065940101_PNU00300IMI.FIT	0231_0065940101_R1S90608DII.FIT	0231_0065940101_R2S90203DII.FIT	0231_0065940101_SCX00000P5S.FIT
0231_0065940101_M2X00000PTH.FIT	0231_0065940101_0MS40401IMI.FIT	0231_0065940101_PNU00300WDX.FIT	0231_0065940101_R1S90709DII.FIT	0231_0065940101_R2S90305DII.FIT	0231_0065940101_SCX00000P6S.FIT
0231_0065940101_0MS00800IMI.FIT	0231_0065940101_0MS00800RFX.FIT	0231_0065940101_0MU00301IMI.FIT	0231_0065940101_R1S90801DII.FIT	0231_0065940101_R2S90406DII.FIT	0231_0065940101_SCX00000P7S.FIT
0231_0065940101_0MS00800THX.FIT	0231_0065940101_0MS00800WDX.FIT	0231_0065940101_0MU00400IMI.FIT	0231_0065940101_R1S90902DII.FIT	0231_0065940101_R2S90507DII.FIT	0231_0065940101_SCX00000P8S.FIT
0231_0065940101_0MS00800WDX.FIT	0231_0065940101_0MS00801IMI.FIT	0231_0065940101_0MU00400WDX.FIT	0231_0065940101_R1S91003DII.FIT	0231_0065940101_R2S90608DII.FIT	0231_0065940101_SCX00000P9S.FIT
0231_0065940101_0MS00801IMI.FIT	0231_0065940101_0MS00801RFX.FIT	0231_0065940101_0MU00401IMI.FIT	0231_0065940101_R1S91104DII.FIT	0231_0065940101_R2S90709DII.FIT	0231_0065940101_SCX00000RAS.ASC
0231_0065940101_0MS01000IMI.FIT	0231_0065940101_0MS01000RFX.FIT	0231_0065940101_0MU00500IMI.FIT	0231_0065940101_R1S91205DII.FIT	0231_0065940101_R2S90801DII.FIT	0231_0065940101_SCX00000ROS.ASC
0231_0065940101_0MS01000WDX.FIT	0231_0065940101_0MS01001IMI.FIT	0231_0065940101_0MU00500WDX.FIT	0231_0065940101_R1S91306DII.FIT	0231_0065940101_R2S90902DII.FIT	0231_0065940101_SCX00000SUM.ASC
0231_0065940101_0MS01001IMI.FIT	0231_0065940101_0MS40000PAX.FIT	0231_0065940101_0MU00501IMI.FIT	0231_0065940101_R1S91408DII.FIT	0231_0065940101_R2S91003DII.FIT	0231_0065940101_SCX00000SUM.SAS
0231_0065940101_0MS40000PAX.FIT	0231_0065940101_0MS40000RFX.FIT	0231_0065940101_0MU00600IMI.FIT	0231_0065940101_R1S91509DII.FIT	0231_0065940101_R2S91105DII.FIT	0231_0065940101_SCX00000TCS.FIT
0231_0065940101_0MS40000RFX.FIT	0231_0065940101_0MS40100IMI.FIT	0231_0065940101_0MU00700IMI.FIT	0231_0065940101_R1S91601DII.FIT	0231_0065940101_R2S91206DII.FIT	0231_0065940101_SCX00000TCX.FIT
0231_0065940101_0MS40100IMI.FIT			0231_0065940101_R1S91702DII.FIT	0231_0065940101_R2S91307DII.FIT	MANIFEST.284964

MOS1

MOS2

OM

pn

RGS1

RGS2


Data analysis

Standard Analysis System (SAS)

The screenshot shows the 'SAS News' section of the XMM-Newton website. At the top, there's a navigation bar with links for 'SCIENCE MISSIONS', 'EUROPEAN SPACE AGENCY', 'SCIENCE & TECHNOLOGY', and 'SIGN IN'. Below the navigation is a banner featuring the 'xmm-newton' logo, a satellite image, and the 'esa' logo.


The main content area has a sidebar on the left with a red header containing 'xm...'. The sidebar includes a list of links such as 'Home / Latest News', 'Conferences & Meetings', 'News', 'General User Support', 'Proposers Info', 'Observers Info', 'Data Analysis', 'Archive, Pipeline & Catalogues', 'Calibration & Background', 'SOC Info', 'About XMM-Newton', 'Image Gallery', 'Publications', and 'Other Links'. To the right of the sidebar, the main content area has a blue header 'SAS NEWS' and a sub-header 'XMM-NEWTON SCIENCE ANALYSIS SOFTWARE'. It displays the current version as '20.0.0' and the release date as 'Dec. 9, 2021'. Below this, there's a paragraph about the Science Analysis Software (SAS) and its purpose. Further down, there's a 'NEWS' section with a bullet point about the release of SAS 20.0.0 binaries on Dec. 9, 2021, and a link to the corresponding release notes. At the bottom of the page is a dark footer bar with a downward arrow icon and the text 'COPYRIGHT 2022 © EUROPEAN SPACE AGENCY. ALL RIGHTS RESERVED.'

2. Data re-processing




* ccf=current calibration files

ODF
(Observation Data Files)



Data Processing



EVT
ccf.cif (Calibration Index Files)

0231_0065940101_AttHk.ds	0231_0065940101_EMOS2_S007_01_Badpixels.ds	0231_0065940101_EPN_S004_02_Badpixels.ds	0231_0065940101_EPN_S004_11_Badpixels.ds
0231_0065940101_EMOS1_S003_01_Badpixels.ds	0231_0065940101_EMOS2_S007_02_Badpixels.ds	0231_0065940101_EPN_S004_03_Badpixels.ds	0231_0065940101_EPN_S004_12_Badpixels.ds
0231_0065940101_EMOS1_S003_02_Badpixels.ds	0231_0065940101_EMOS2_S007_03_Badpixels.ds	0231_0065940101_EPN_S004_04_Badpixels.ds	0231_0065940101_EPN_S004_ImagingEvts.ds
0231_0065940101_EMOS1_S003_03_Badpixels.ds	0231_0065940101_EMOS2_S007_04_Badpixels.ds	0231_0065940101_EPN_S004_05_Badpixels.ds	ccf.cif
0231_0065940101_EMOS1_S003_04_Badpixels.ds	0231_0065940101_EMOS2_S007_05_Badpixels.ds	0231_0065940101_EPN_S004_06_Badpixels.ds	m1.evt
0231_0065940101_EMOS1_S003_05_Badpixels.ds	0231_0065940101_EMOS2_S007_06_Badpixels.ds	0231_0065940101_EPN_S004_07_Badpixels.ds	m2.evt
0231_0065940101_EMOS1_S003_06_Badpixels.ds	0231_0065940101_EMOS2_S007_07_Badpixels.ds	0231_0065940101_EPN_S004_08_Badpixels.ds	pn.evt
0231_0065940101_EMOS1_S003_07_Badpixels.ds	0231_0065940101_EMOS2_S007_ImagingEvts.ds	0231_0065940101_EPN_S004_09_Badpixels.ds	
0231_0065940101_EMOS1_S003_ImagingEvts.ds	0231_0065940101_EPN_S004_01_Badpixels.ds	0231_0065940101_EPN_S004_10_Badpixels.ds	



odtingest



emproc (for EPIC-MOS)



epproc (for EPIC-pn)


Event files



Contains all the CCF
necessary for your
observation

Data reduction

SAS threads



IMAGE

SPECTRUM

LIGHT CURVE

fv pn.evt

#rows=#counts

fv: Summary of pn.evt in /blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn/

Index	Extension	Type	Dimension	View				
0	Primary	Image	0	Header	Image	Table		
1	EVENTS	Binary	15 cols X 131297 rows	Header	Hist	Plot	All	Select
2	OFFSETS	Binary	3 cols X 14 rows	Header	Hist	Plot	All	Select
3	EXPOSU01	Binary	2 cols X 95904 rows	Header	Hist	Plot	All	Select
4	BADPIX01	Binary	5 cols X 72 rows	Header	Hist	Plot	All	Select
5	DLIMAP01	Binary	3 cols X 200 rows	Header	Hist	Plot	All	Select
6	HKAUX01	Binary	2 cols X 4789 rows	Header	Hist	Plot	All	Select
7	EXPOSU02	Binary	2 cols X 95901 rows	Header	Hist	Plot	All	Select
8	BADPIX02	Binary	5 cols X 68 rows	Header	Hist	Plot	All	Select
9	DLIMAP02	Binary	3 cols X 200 rows	Header	Hist	Plot	All	Select

X,Y -> image
Time -> light curve
Energy -> spectrum

fv pn.evt

fv: Binary Table of pn.evt[1] in /blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn/

	TIME	RAWX	RAWY	DETX	DETY	X	Y	PHA	PI	FLAG	PATTERN	PAT_ID	PAT_SEQ	CCDNR	TIME_RAW
Select	D	I	I	I	I	J	J	I	I	J	B	I	B	B	D
All	s	pixel	pixel	0.05 arcsec	0.05 arcsec	0.05 arcsec	0.05 arcsec	channel	eV						s
Invert	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify
1	1.910934147154E+08	23	24	-4118	13485	18376	16096	24	211	0	1	5121	0	1	1.910934146937E+08
2	1.910934172390E+08	41	84	-5619	8551	18939	21222	158	835	0	0	0	0	1	1.910934172615E+08
3	1.910934173916E+08	7	47	-2815	11608	20313	17308	21	209	0	4	5121	0	1	1.910934174082E+08
4	1.910934174143E+08	13	47	-3294	11601	19875	17503	33	279	0	2	5122	0	1	1.910934174082E+08
5	1.910934173875E+08	58	64	-7048	10182	16984	20286	2635	13720	0	0	0	0	1	1.910934174082E+08
6	1.910934184187E+08	14	119	-3355	5690	22147	22960	111	585	0	0	0	0	1	1.910934184353E+08
7	1.910934196784E+08	34	125	-5034	5146	20818	24122	2918	24348	0	2	5121	0	1	1.910934196825E+08
8	1.910934197620E+08	45	14	-5919	14325	16389	16033	30	400	1	78	1	1	1	1.910934197559E+08
9	1.910934199950E+08	10	115	-3048	6034	22294	22523	30	158	0	0	0	0	1	1.910934199760E+08
10	1.910934209479E+08	64	146	-7536	3436	19192	26679	377	2097	4	0	0	0	1	1.910934209297E+08
11	1.910934212498E+08	11	42	-3096	11994	19902	17064	965	4788	0	0	0	0	1	1.910934212232E+08
12	1.910934246840E+08	39	13	-5436	14441	16788	15736	39	192	32	0	0	0	1	1.910934246713E+08
13	1.910934246531E+08	39	17	-5405	14090	16954	16047	31	692	1	96	1	1	1	1.910934246713E+08
14	1.910934253051E+08	4	111	-2520	6289	22679	22081	31	160	0	0	0	0	1	1.910934253316E+08
15	1.910934259405E+08	19	173	-3791	1226	23505	27235	40	297	8	0	0	0	1	1.910934259185E+08
16	1.910934283720E+08	11	122	-3167	5397	22436	23155	255	1330	0	0	0	0	1	1.910934283396E+08
17	1.910934289507E+08	49	146	-6305	3404	20336	26223	2870	15263	0	0	0	0	1	1.910934289265E+08
18	1.910934291953E+08	6	138	-2736	4076	23352	24200	866	6695	0	1	5121	0	1	1.910934292200E+08
19	1.910934303410E+08	10	161	-3039	2187	23818	26056	28	155	0	0	0	0	1	1.910934303205E+08
20	1.910934320374E+08	23	146	-4097	3414	22362	25344	42	225	0	0	0	0	1	1.910934320078E+08
21	1.910934323060E+08	43	175	-5762	1046	21764	28177	2745	19547	0	7	5121	1	1	1.910934323013E+08
22	1.910934341115E+08	4	16	-2581	14133	19533	14895	26	232	0	3	5121	1	1	1.910934341354E+08
23	1.910934341378E+08	4	21	-2555	13784	19695	15205	34	160	0	0	0	5	1	1.910934341354E+08
24	1.910934362451E+08	61	13	-7263	14438	15110	16459	159	1076	1	2	1	0	1	1.910934362630E+08
25	1.910934363497E+08	54	140	-6705	3949	19754	25880	62	343	0	0	0	0	1	1.910934363364E+08
26	1.910934365385E+08	43	184	-5761	328	22048	28836	27	158	0	0	0	0	1	1.910934365565E+08

Go to:

Edit cell:



3. Filtering against high background

EPIC particle induced background

(above a few keV)

Internal 'quiescent' component

high energy particles interacting with the structure surrounding the detectors and the detectors themselves




3. Filtering against high background

EPIC particle induced background


(above a few keV)

External 'flaring' component

strong and rapid variability; currently attributed to soft protons ($E_p <$ a few 100 keV) likely organized in clouds populating the Earth's magneto-sphere (above a few keV)





MOS2 observation
badly affected by soft proton flares



MOS1 light curve
badly affected by soft proton flares

Internal 'quiescent' component

high energy particles interacting with the structure surrounding the detectors and the detectors themselves (below 300 eV)



Creation of a light curve above 10 keV

How to filter an EPIC event list for periods of high background *flaring* activity

Extract a **single event (i.e. pattern zero only)**, high energy light curve, from the event file to identify intervals of flaring particle background:

pn

```
evselect table=pn.evt energycolumn=PI expression='#XMMEA_EP && (PI>10000&&PI<12000) && (PATTERN==0)'  
withrateset=yes rateset="lcurve_sup10.lc" timebinsize=100 maketimecolumn=yes makeratecolumn=yes
```

Notice that the EPIC-pn energy range selected for producing the background light curve includes events only up to 12 keV. The reason for this is to avoid hot pixels being miss-identified as very high energy events.

maketimecolumn=If true, include a time column in the FITS table when creating a time series;

makeratecolumn=If true, produces a lightcurve containing a RATE, rather than a COUNTS column. An ERROR column is also produced.

Creation of a light curve above 10 keV

How to filter an EPIC event list for periods of high background *flaring* activity

MOS 1

```
evselect table=m1.evt energycolumn=PI expression='#XMMEA_EM && (PI>10000) &&(PATTERN==0)' withrateset=yes
rateset="lcurve_sup10.lc" timebinsize=100 maketimecolumn=yes makeratecolumn=yes
```

MOS 2

```
evselect table=m2.evt energycolumn=PI expression='#XMMEA_EM && (PI>10000) &&(PATTERN==0)' withrateset=yes
rateset="lcurve_sup10.lc" timebinsize=100 maketimecolumn=yes makeratecolumn=yes
```

But what are event patterns?

Table 7: List of EPIC event patterns

Camera	Mode	X-ray generated pattern			
		singles	doubles	triples	quadruples
MOS	imaging	0	1-4	5-8	9-12
pn	imaging	0	1-4	5-8	9-12

Events that are recommended
for spectral analysis

single event X
double pattern	. . .	X X x .	. . X x X
triple pattern x X X x .	. . X x .	. . x X
quadruple pattern m x x m
 x X X x .	. . X x .	. . X x .	. . x X

Figure 13: List of valid EPIC-pn patterns (cf. figure 12). Here " ." marks a pixel without an event above threshold, "X" is the pixel with the maximum charge ("main pixel"), "x" is the pixel with a non-maximum charge, "m" is the pixel with the minimum charge. These 13 figures refer to the SAS PATTERN codes 0 (singles), 1-4 (doubles), 5-8 (triples) and 9-12 (quadruples), respectively. The RAWX co-ordinate is running rightward and the RAWY co-ordinate running upward.

> lcurve

To visualize the produced lightcurve

```
torresi@login06:/blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn [torresi@login06]pn>lcurve

lcurve 1.0 (xronos6.0)

Number of time series for this task[1]
Ser. 1 filename +options (or @file of filenames +options)[lcurve_sup10.lc]
  Series 1 file    1:lcurve_sup10.lc

Selected FITS extensions: 1 - RATE TABLE;

Source ..... Start Time (d) .... 13025 17:30:12.548
FITS Extension .... 1 - `RATE` Stop Time (d) .... 13025 19:27:28.859
No. of Rows ..... 71 Bin Time (s) .... 100.0
Right Ascension ... Internal time sys.. Converted to TJD
Declination ..... Experiment ..... XMM      EPN
Filter ..... Medium
Corrections applied: Vignetting - No ; Deadtime - No ; Bkgd - No ; Clock - Yes

Selected Columns: 3- Time; 1- Y-axis; 2- Y-error;

File contains binned data.

Name of the window file ('-' for default window)[-]

Expected Start ... 13025.72931189543 (days) 17:30:12:548 (h:m:s:ms)
Expected Stop .... 13025.81075068651 (days) 19:27:28:859 (h:m:s:ms)

Minimum Newbin Time 100.00000 (s)
for Maximum Newbin No.. 71


Default Newbin Time is: 100.00000 (s) (to have 1 Intv. of 71 Newbins)
Type INDEF to accept the default value

Newbin Time or negative rebinning[100]



Newbin Time ..... 100.00000 (s)
Maximum Newbin No. 71

Default Newbins per Interval are: 71
Giving 1 Interval of 71 Newbins
Type INDEF to accept the default value

Number of Newbins/Interval[109] 71
Maximum of 1 Intvs. with 71 Newbins of 100.000 (s)
Name of output file[test.flc]
Do you want to plot your results?[yes]
```




Start Time 13025 17:31:02:548 Stop Time 13025 19:27:42:548



Possible error! Disconnect from login06 and reconnect

PGPLOT /xw: cannot connect to X server [localhost:12.0]
To plot vs. Time (s), please enter
PGPLOT file/type:

4. Selection of GOOD TIME INTERVALS (GTI)



```
tabgtigen table=lcurve_sup10_lc gtiset=good_bkg.gti expression='RATE<0.7'
```

5. Generation of the cleaned event file

pn

```
evselect table=pn.evt expression='#XMMEA_EP && (PI>150) && (GTI(good_bkg.gti,TIME))' withfilteredset=yes
keepfilteroutput=yes filteredset=pn_new.evt updateexposure=yes cleandss=yes writedss=yes
```

filteredset= the name of the file to which the filtered event list is to be written. If this parameter is set, then **keepfilteroutput** is automatically set to true.

updateexposure= update exposure information in event lists (keywords LIVETIME, LIVETInn, ONTIME, and ONTIMEnn) and in spectrum files (keyword EXPOSURE).

cleandss= controls the use of data subspace cleaning, which deletes components from the data subspace which select no events from the event list.

writess= controls the writing of data subspace information to the output data files.

5. Generation of the cleaned event file

pn

```
evselect table=pn.evt expression='#XMMEA_EP && (PI>150) && (GTI(good_bkg.gti,TIME))' withfilteredset=yes  
keepfilteroutput=yes filteredset=pn_new.evt updateexposure=yes cleandss=yes writedss=yes
```

fv: Binary Table of pn.evt[1] in /blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn/

	TIME	RAWX	RAWY	DETX	DETY	X	Y	PHA	PI	FLAG	PATTERN
Select	D	I	I	I	I	J	J	I	I	J	B
All	s	pixel	pixel	0.05 arcsec	0.05 arcsec	0.05 arcsec	0.05 arcsec	channel	EV		
Invert	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify	Modify
1	1.910949802434E+08	50	110	7331	-8630	37610	31914	30	151	0	0
2	1.910936738810E+08	28	142	-9935	3770	16856	27317	24	151	0	0
3	1.910987199062E+08	20	183	-11361	-2593	18051	33727	24	151	0	0
4	1.910968628259E+08	35	24	11477	-15702	44206	36781	30	151	65536	0
5	1.910990361903E+08	29	187	-10571	-2284	18655	33132	24	151	0	0
6	1.910961221517E+08	62	98	-7817	-9592	24065	38764	31	151	2097160	0
7	1.910970038855E+08	1	133	3105	4529	28542	21483	26	151	4	0
8	1.910974374096E+08	62	45	-7897	-13947	25707	42799	33	151	2097160	0
9	1.910945815748E+08	48	181	-11601	516	16606	30964	22	151	0	0
10	1.910967549282E+08	15	169	-14269	1567	13740	31049	25	151	0	0
11	1.910971991752E+08	56	151	3922	3037	29881	22532	26	151	0	0
12	1.910989785826E+08	35	162	5695	2135	31866	22663	24	151	0	0
13	1.910968081495E+08	57	186	-13624	-2308	15859	34356	22	151	0	0
14	1.910990163098E+08	48	14	-6206	14311	16131	16159	30	151	0	0
15	1.910954398068E+08	46	116	-16806	5925	9691	28042	27	151	0	0
16	1.910962318820E+08	21	160	-5831	-4511	23889	33312	28	151	0	0
17	1.910999689040E+08	1	167	-2130	-3895	27049	31288	31	151	4	0
18	1.910993828217E+08	57	134	-17669	4455	9477	29733	26	151	0	0
19	1.910989936606E+08	60	164	-12546	1956	15170	30012	22	151	0	0
20	1.910960863435E+08	41	151	-14993	-5262	15764	37611	27	151	0	0
21	1.910968742872E+08	43	185	12176	-2456	39631	24330	24	151	0	0
22	1.910997870684E+08	51	199	2000	-1237	29798	27218	30	151	0	0
23	1.910989410890E+08	44	177	-14728	-3097	15155	35516	22	151	0	0
24	1.910960325198E+08	37	194	-9900	-1697	19041	32328	24	151	0	0
25	1.910967498674E+08	33	154	11287	-5002	39817	27021	26	151	0	0
26	1.910957214857E+08	57	167	-13697	-3921	16427	35868	23	151	0	0

Go to:

Edit cell: 151

5. Generation of the cleaned event file

MOS 1

```
evselect table=m1.evt expression='#XMMEA_EM && (PI > 150) && (GTI(good_bkg.gti,TIME))' withfilteredset=yes  
keepfilteroutput=yes filteredset=mos1_new.evt updateexposure=yes cleandss=yes writedss=yes
```

MOS 2

```
evselect table=m2.evt expression='#XMMEA_EM && (PI > 150) && (GTI(good_bkg.gti,TIME))' withfilteredset=yes  
keepfilteroutput=yes filteredset=mos2_new.evt updateexposure=yes cleandss=yes writedss=yes
```

Raw event file (pn.evt)

fv: Summary of pn.evt in /blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn/

File	Edit	Tools	Help						
Index	Extension	Type	Dimension	View					
0	Primary	Image	0	Header	Image	Table			
1	EVENTS	Binary	15 cols X 131297 rows	Header	Hist	Plot	All	Select	
2	OFFSETS	Binary	3 cols X 14 rows	Header	Hist	Plot	All	Select	
3	EXPOSU01	Binary	2 cols X 95904 rows	Header	Hist	Plot	All	Select	
4	BADPIX01	Binary	5 cols X 72 rows	Header	Hist	Plot	All	Select	
5	DLIMAP01	Binary	3 cols X 200 rows	Header	Hist	Plot	All	Select	
6	HKAUX01	Binary	2 cols X 4789 rows	Header	Hist	Plot	All	Select	
7	EXPOSU02	Binary	2 cols X 95901 rows	Header	Hist	Plot	All	Select	
8	BADPIX02	Binary	5 cols X 68 rows	Header	Hist	Plot	All	Select	
9	DLIMAP02	Binary	3 cols X 200 rows	Header	Hist	Plot	All	Select	



Cleaned event file (pn_new.evt)

fv: Summary of pn_new.evt in /blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn/

Is	Help								
Extension	Type	Dimension	View						
0	Primary	Image	0	Header	Image	Table			
1	EVENTS	Binary	15 cols X 125977 rows	Header	Hist	Plot	All	Select	
2	OFFSETS	Binary	3 cols X 14 rows	Header	Hist	Plot	All	Select	
3	EXPOSU01	Binary	2 cols X 95904 rows	Header	Hist	Plot	All	Select	
4	BADPIX01	Binary	5 cols X 72 rows	Header	Hist	Plot	All	Select	
5	DLIMAP01	Binary	3 cols X 200 rows	Header	Hist	Plot	All	Select	
6	HKAUX01	Binary	2 cols X 4789 rows	Header	Hist	Plot	All	Select	
7	EXPOSU02	Binary	2 cols X 95901 rows	Header	Hist	Plot	All	Select	
8	BADPIX02	Binary	5 cols X 68 rows	Header	Hist	Plot	All	Select	
9	DLIMAP02	Binary	3 cols X 200 rows	Header	Hist	Plot	All	Select	

SAOImage ds9

File Edit View Frame Bin Zoom Scale Color Region WCS Analysis Help

File pn_new.evt[EVENTS]

Object 3C33


Value 0

FK5 α 1:09:04.7108 δ +13:30:54.907

Physical x 24196.9 y 39855.3


Image x 276.334 y 519.465

Frame 1 x 0.98743 y 0 .



file edit view frame bin zoom scale color region wcs analysis help


zoom in zoom out zoom fit zoom 1/8 zoom 1/4 zoom 1/2 zoom 1 zoom 2 zoom 4 zoom 8



Full frame

0.1 0.4 0.9 1.8 3.8 7.7 15.5 31.1 62.1

Zoomed image with radio (VLA 5GHz) contours overlaid



6. Source extraction region


Display the cleaned pn* image with **ds9**

ds9 pn_new.evt &

> scale log
> bin (block 2, 4, ...)

> Region
> save region
> file format ‘ds9’
> coordinates ‘physical’
> **source.reg**

<http://ds9.si.edu/doc/user/binning/index.html>





circle(27720.501,27069.104,700.00011)

* or MOS1, MOS2



Fraction encircled energy

Fraction of photons contained within a certain angular radius



on-axis — point-like source

Extraction radius (arcsec)	Camera		
	MOS-1	MOS2	EPN pointed
15	0.68	0.69	0.71
30	0.83	0.83	0.88
45	0.89	0.89	0.93
60	0.92	0.93	0.95



Each telescope has its own on-axis PSF;

the core of XMM on-axis PSF is **narrow** and **varies little** in the 0.1-6 keV band (major XMM strong point!)

see also Dadina's lesson

7. Background extraction region


Display the cleaned pn image with **ds9**

```
ds9 pn_new.evt &
```


```
> scale log  
> bin (block 2, 4, ...)
```

```
> Region  
> save region  
> file format 'ds9'  
> coordinates 'physical'  
> back.reg
```

```
circle(25864.501,24541.104,700.00011)
```




EPIC Science Modes






Imaging modes

Timing Mode




MOS






Partial Window





pn



RAWX






PILEUP

<https://www.cosmos.esa.int/web/xmm-newton/sas-thread-epatplot>

Arrival of two or more independent photons at nearby pixels that are erroneously read as one single event (whose energy is the sum of the energies of the individual photons) [Jethwa et al. \(2015\)](#)



single event	. X .			
	. . .			
	. . .			
	. . .			
	. . .			
double pattern	. . X X x .	. . X . .	. x X . .
 X

triple pattern	. x X X x .	. . X x .	. x X . .
 X X . .

quadruple pattern	. m X X m .	. . X x .	. x X . .
	. x X X x .	. . X x .	. x X . .
 X m .	. m X . .

Figure 13: List of valid EPIC-pn patterns (cf. figure 12). Here "." marks a pixel without an event above threshold, "X" is the pixel with the maximum charge ("main pixel"), "x" is the pixel with a non-maximum charge, "m" is the pixel with the minimum charge. These 13 figures refer to the SAS PATTERN codes 0 (singles), 1-4 (doubles), 5-8 (triples) and 9-12 (quadruples), respectively. The RAWX co-ordinate is running rightward and the RAWY co-ordinate running upward.

- Photon loss
 - Energy distortion
 - Pattern migration

8. Check for the presence of pileup

Diagnostic tool for pileup: **epatplot**

```
evselect table=pn_new.evt withfilteredset=yes filteredset=pnf.evt keepfilteroutput=yes expression="((X,Y) IN circle (25910.5,25870.5,400))"
```

```
epatplot set=pf.evt device="/CPS" plotfile="pnf_pat.ps" or device="/GIF" plotfile="pnf_pat.gif"
```


```
[torresi@login06]pn>export SAS_CCF='/blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/evt/ccf.cif'
[torresi@login06]pn>epatplot set=pf.evt device="/CPS" plotfile="pnf_pat.ps"
epatplot:- Executing (routine): epatplot set=pf.evt modifyinset=yes sigma=3 withflag=yes xaxisadu=no device=/CPS o
utdir=./ plotfile=pf_pat.ps useplotfile=yes withqdp=no withdetxy=no withsrcxy=yes outmaskname=flag0_map_##.dat wit
houtputmask=no backgroundset=bkg_events.fits withbackgroundset=no backscal=1 usermode=0 withusermode=no userrawy=19
0 withuserrawy=no ccdlimits='1 64 1 200 1 12' plot xrange='0 0' plot yrange='0 0' pileupnumberenergyrange='500 2000'
-w 1 -V 4
epatplot:- epatplot (epatplot-1.22) [xmmsas_20211130_0941-20.0.0] started: 2022-11-16T10:57:22.000
epatplot:- epatplot 1.22 is running...
epatplot:-
epatplot:-      s      d      t      q
epatplot:-  0.6375  0.3228  0.0176  0.0220
epatplot:-
epatplot:- 0.5-2.0 keV observed-to-model fractions:
epatplot:- s: 0.955 +/- 0.094   d: 1.168 +/- 0.156
epatplot:-
epatplot:- PostScript output written to file pnf_pat.ps
epatplot:- epatplot (epatplot-1.22) [xmmsas_20211130_0941-20.0.0] ended: 2022-11-16T10:57:23.000
[torresi@login06]pn>gv pnf_pat.ps&
[11:12:57]
```

Epatplot compares the ratios of single- and double-events with standard values

gv pnf_pat.ps

evince pnf_pat.ps

gimp pnf_pat.gif





The source is not piled up!

Table 3: Basic numbers for the science modes of EPIC


MOS (central CCD; pixels) [1 pixel = 1.1"]	Time resolution	Live time ¹ [%]	Max. count rate ² diffuse ³ (total) [s ⁻¹]	Max. count rate ² (flux) point source [s ⁻¹] ([mCrab] ⁴)
Full frame (600×600)	2.6 s	100.0	150	0.50 (0.17)
Large window (300×300)	0.9 s	99.5	110	1.5 (0.49)
Small window (100×100)	0.3 s	97.5	37	4.5 (1.53)
Timing uncompressed (100×600)	1.75 ms	100.0	N/A	100 (35)

pn (array or 1 CCD; pixels) [1 pixel = 4.1"]	Time resolution	Live time ¹ [%]	Max. count rate ² diffuse ³ (total) [s ⁻¹]	Max. count rate ² (flux) point source [s ⁻¹] ([mCrab] ⁴)
Full frame ⁵ (376×384)	73.4 ms	99.9	1000(total)	2 (0.23)
Extended full frame ^{5,6} (376×384)	199.1 ms	100.0	370	0.7 (0.09)
Large window (198×384)	47.7 ms	94.9	1500	3 (0.35)
Small window (63×64)	5.7 ms	71.0	12000	25 (3.25)
Timing (64×200)	0.03 ms	99.5	N/A	800 (85)
Burst (64×180)	7 μs	3.0	N/A	60000 (6300)





While, in this case, the source **is** piled up



9. SPECTRUM EXTRACTION

pn

```
evselect table=pn_new.evt withspectrumset=yes spectrumset=source_spectrum.fits energycolumn=PI spectralbinsize=5
withspecranges=yes specchannelmin=0 specchannelmax=20479 expression='(FLAG==0) && (PATTERN<=4) && ((X,Y) IN circle
(27874.528,26645.58,699.9999))'
```

source.reg

spectralbinsize= binning factor for spectral creation (size of each bin in instrumental eV)

specchannelmin= the minimum channel number to consider for spectrum creation

specchannelmax= the maximum channel number to consider for spectrum creation

9. SPECTRUM EXTRACTION

MOS 1

```
evselect table=mos1_new.evt withspectrumset=yes spectrumset=source_spectrum.fits energycolumn=PI spectralbinsize=15
withspecranges=yes specchannelmin=0 specchannelmax=11999 expression='(FLAG==0) && (PATTERN<=12) && ((X,Y) IN circle
(28090.5,24221.5,775.48791))'
```

MOS 2

```
evselect table=mos2_new.evt withspectrumset=yes spectrumset=source_spectrum.fits energycolumn=PI spectralbinsize=15
withspecranges=yes specchannelmin=0 specchannelmax=11999 expression='(FLAG==0) && (PATTERN<=12) && ((X,Y) IN circle
(28090.5,24221.5,775.48791))'
```

10. BACKGROUND EXTRACTION

pn

```
evselect table=pn_new.evt withspectrumset=yes spectrumset=back_spectrum.fits energycolumn=PI spectralbinsize=5
withspecranges=yes specchannelmin=0 specchannelmax=20479 expression='(FLAG==0) && (PATTERN<=4) && ((X,Y) IN circle
(25864.501,24541.104,700.00011))'
```

back.reg

10. BACKGROUND EXTRACTION

MOS 1

```
evselect table=mos1_new.evt withspectrumset=yes spectrumset=back_spectrum.fits energycolumn=PI spectralbinsize=15
withspecranges=yes specchannelmin=0 specchannelmax=11999 expression='(FLAG==0) && (PATTERN<=12) && ((X,Y) IN circle
(25864.501,24541.104,700.00011))'
```

MOS 2

```
evselect table=mos2_new.evt withspectrumset=yes spectrumset=back_spectrum.fits energycolumn=PI spectralbinsize=15
withspecranges=yes specchannelmin=0 specchannelmax=11999 expression='(FLAG==0) && (PATTERN<=12) && ((X,Y) IN circle
(25864.501,24541.104,700.00011))'
```

11. BACKSCALE

The BACKSCALE task calculates the area of a source region used to make a spectral file.

This task takes into account any bad pixels or chip gaps and writes the result into the BACKSCAL keyword of the SPECTRUM table

The final value is:

AREA= GEOMETRIC AREA-CCD GAPS-BAD PIXELS

pn

```
backscale spectrumset=source_spectrum.fits badpixlocation=pn_new.evt
backscale spectrumset=back_spectrum.fits badpixlocation=pn_new.evt
```

MOS 1

```
backscale spectrumset=source_spectrum.fits badpixlocation=mos1_new.evt
backscale spectrumset=back_spectrum.fits badpixlocation=mos1_new.evt
```

MOS 2

```
backscale spectrumset=source_spectrum.fits badpixlocation=mos2_new.evt
backscale spectrumset=back_spectrum.fits badpixlocation=mos2_new.evt
```

11. BACKSCALE

The BACKSCALE task calculates the area of a source region used to make a spectral file.

This task takes into account any bad pixels or chip gaps and writes the result into the BACKSCAL keyword of the SPECTRUM table

fv: Header of source_spectrum.fits[1] in /blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn/ - □ ×

File Edit Tools Help

Search for: BACKSCAL Case sensitive? No

```
RESPFILE= 'NONE'          / Name of response file
DSTYP1 = 'CCDNR'          / data subspace descriptor: name
DSTYP2 = 'FLAG'            / data subspace descriptor: name
DSTYP3 = 'P1'              / data subspace descriptor: name
DSUNI3 = 'eV'              / data subspace descriptor: units
DSTYP4 = 'TIME'            / data subspace descriptor: name
DSUNI4 = 's'                / data subspace descriptor: units
DSTYP5 = 'FLAG'            / data subspace descriptor: name
DSTYP6 = 'PATTERN'         / data subspace descriptor: name
DSTYP7 = 'POS(X,Y)'        / data subspace descriptor: name
DSVAL1 = '1'                / data subspace descriptor: value
DSFORM2 = 'X'              / data subspace descriptor: data type
DSVAL2 = 'b00xx00000x0xxxxxxxxxxxxxx' / data subspace descriptor: value
DSVAL3 = '(150:'           / data subspace descriptor: value
DSVAL4 = 'TABLE'            / data subspace descriptor: value
DSREF4 = ':GTI00004'         / data subspace descriptor: reference
DSVAL5 = '0'                / data subspace descriptor: value
DSVAL6 = ':4'               / data subspace descriptor: value
DSVAL7 = 'TABLE'            / data subspace descriptor: value
DSREF7 = ':REG00107'         / data subspace descriptor: reference
2DSVAL1 = '2'                / data subspace descriptor: value
2DSREF4 = ':GTI00104'         / data subspace descriptor: reference
3DSVAL1 = '3'                / data subspace descriptor: value
3DSREF4 = ':GTI00204'         / data subspace descriptor: reference
4DSVAL1 = '4'                / data subspace descriptor: value
4DSREF4 = ':GTI00304'         / data subspace descriptor: reference
5DSVAL1 = '5'                / data subspace descriptor: value
5DSREF4 = ':GTI00404'         / data subspace descriptor: reference
6DSVAL1 = '6'                / data subspace descriptor: value
6DSREF4 = ':GTI00504'         / data subspace descriptor: reference
7DSVAL1 = '7'                / data subspace descriptor: value
7DSREF4 = ':GTI00604'         / data subspace descriptor: reference
8DSVAL1 = '8'                / data subspace descriptor: value
8DSREF4 = ':GTI00704'         / data subspace descriptor: reference
9DSVAL1 = '9'                / data subspace descriptor: value
9DSREF4 = ':GTI00804'         / data subspace descriptor: reference
10DSVAL1= '10'               / data subspace descriptor: value
10DSREF4= ':GTI00904'         / data subspace descriptor: reference
11DSVAL1= '11'               / data subspace descriptor: value
11DSREF4= ':GTI01004'         / data subspace descriptor: reference
12DSVAL1= '12'               / data subspace descriptor: value
12DSREF4= ':GTI01104'         / data subspace descriptor: reference
MTYPE1 = 'POS'              / DM Keyword: Descriptor name
MFORM1 = 'X,Y'              / DM Keyword: Descriptor name
EXPOSURE= 6.27215711148456E+03 / Weighted live time of CCDs in the extraction re
BACKSCAL= 1542180 / Scaling factor for background
TLMIN1 = 0
TLMAX1 = 4095
END
```

BACKSCAL= 1542180 / Scaling factor for background

Source

fv: Header of back_spectrum.fits[1] in /blasco/users/torresi/LAB-X/Fall_2022/3C33_LABX/pn/ - □ ×

File Edit Tools Help

Search for: Case sensitive? No

```
DSVAL2 = 'b00xx00000x0xxxxxxxxxxxxxx' / data subspace descriptor: value
DSVAL3 = '(150:'           / data subspace descriptor: value
DSVAL4 = 'TABLE'            / data subspace descriptor: value
DSREF4 = ':GTI00004'         / data subspace descriptor: reference
DSVAL5 = '0'                / data subspace descriptor: value
DSVAL6 = ':4'               / data subspace descriptor: value
DSVAL7 = 'TABLE'            / data subspace descriptor: value
DSREF7 = ':REG00107'         / data subspace descriptor: reference
2DSVAL1 = '2'                / data subspace descriptor: value
2DSREF4 = ':GTI00104'         / data subspace descriptor: reference
3DSVAL1 = '3'                / data subspace descriptor: value
3DSREF4 = ':GTI00204'         / data subspace descriptor: reference
4DSVAL1 = '4'                / data subspace descriptor: value
4DSREF4 = ':GTI00304'         / data subspace descriptor: reference
5DSVAL1 = '5'                / data subspace descriptor: value
5DSREF4 = ':GTI00404'         / data subspace descriptor: reference
6DSVAL1 = '6'                / data subspace descriptor: value
6DSREF4 = ':GTI00504'         / data subspace descriptor: reference
7DSVAL1 = '7'                / data subspace descriptor: value
7DSREF4 = ':GTI00604'         / data subspace descriptor: reference
8DSVAL1 = '8'                / data subspace descriptor: value
8DSREF4 = ':GTI00704'         / data subspace descriptor: reference
9DSVAL1 = '9'                / data subspace descriptor: value
9DSREF4 = ':GTI00804'         / data subspace descriptor: reference
10DSVAL1= '10'               / data subspace descriptor: value
10DSREF4= ':GTI00904'         / data subspace descriptor: reference
11DSVAL1= '11'               / data subspace descriptor: value
11DSREF4= ':GTI01004'         / data subspace descriptor: reference
12DSVAL1= '12'               / data subspace descriptor: value
12DSREF4= ':GTI01104'         / data subspace descriptor: reference
MTYPE1 = 'POS'              / DM Keyword: Descriptor name
MFORM1 = 'X,Y'              / DM Keyword: Descriptor name
EXPOSURE= 6.27215711148456E+03 / Weighted live time of CCDs in the extraction re
BACKSCAL= 1540700 / Scaling factor for background
TLMIN1 = 0
TLMAX1 = 4095
END
```

BACKSCAL= 1540700 / Scaling factor for background

Background

12. Redistribution Matrix File (RMF)


```
rmfgen spectrumset=source_spectrum.fits rmfset=pn.rmf
```

The RMF associates to each instrument channel (I) the appropriate photon energy (E)

fv: Binary Table of `ftm0830p3759.rmf[2]` in /hc

	CHANNEL	E_MIN	E_MAX
Select	1E	1E	1E
All	channel	keV	keV
1	1.000000E+00	1.460000E-03	1.460000E-02
2	2.000000E+00	1.460000E-02	2.920000E-02
3	3.000000E+00	2.920000E-02	4.380000E-02
4	4.000000E+00	4.380000E-02	5.840000E-02
5	5.000000E+00	5.840000E-02	7.300000E-02
6	6.000000E+00	7.300000E-02	8.760000E-02
7	7.000000E+00	8.760000E-02	1.022000E-01
8	8.000000E+00	1.022000E-01	1.168000E-01
9	9.000000E+00	1.168000E-01	1.314000E-01
10	1.000000E+01	1.314000E-01	1.460000E-01
11	1.100000E+01	1.460000E-01	1.606000E-01
12	1.200000E+01	1.606000E-01	1.752000E-01
13	1.300000E+01	1.752000E-01	1.898000E-01
14	1.400000E+01	1.898000E-01	2.044000E-01
15	1.500000E+01	2.044000E-01	2.190000E-01
16	1.600000E+01	2.190000E-01	2.336000E-01
17	1.700000E+01	2.336000E-01	2.482000E-01
18	1.800000E+01	2.482000E-01	2.628000E-01
19	1.900000E+01	2.628000E-01	2.774000E-01
20	2.000000E+01	2.774000E-01	2.920000E-01

Go to: Edit cell: 0.219



13. Ancillary Response File (ARF)

```
arfgen spectrumset=source_spectrum.fits arfset=pn.arf withrmfset=yes rmfset=pn.rmf  
badpixlocation=pn_new.evt detmaptype=psf (for point sources; for extended sources -> flat)
```



The ARF includes information on the **effective area, filter transmission and any additional energy-dependent efficiencies**, i.e. the efficiency of the instrument in revealing photons

fv: Binary Table of ftm0830p3759.arf[1] in /ho

	ENERG_LO	ENERG_HI	SPECRESP
Select	1E	1E	1E
	keV	keV	cm**2
1	2.200000E-01	2.300000E-01	9.414584E+01
2	2.300000E-01	2.400000E-01	1.119709E+02
3	2.400000E-01	2.500000E-01	1.309653E+02
4	2.500000E-01	2.600000E-01	1.518642E+02
5	2.600000E-01	2.700000E-01	1.716482E+02
6	2.700000E-01	2.800000E-01	1.922011E+02
7	2.800000E-01	2.900000E-01	4.741680E+01
8	2.900000E-01	3.000000E-01	2.284590E+00
9	3.000000E-01	3.100000E-01	5.144246E+00
10	3.100000E-01	3.200000E-01	1.563580E+01
11	3.200000E-01	3.300000E-01	2.251595E+01
12	3.300000E-01	3.400000E-01	3.011008E+01
13	3.400000E-01	3.500000E-01	3.743014E+01
14	3.500000E-01	3.600000E-01	4.385400E+01
15	3.600000E-01	3.700000E-01	4.954287E+01
16	3.700000E-01	3.800000E-01	5.625348E+01
17	3.800000E-01	3.900000E-01	6.431229E+01
18	3.900000E-01	4.000000E-01	7.319862E+01
19	4.000000E-01	4.100000E-01	7.713167E+01
20	4.100000E-01	4.200000E-01	8.444775E+01




Graph coordinates: (4.0175, 610.543)
Image pixel: (X, X)
Pixel value: X ()

Go to: Edit cell: 0.42



** arfgen: fatal error (NoCifSpecified), `/prod_oasbo/sas/xmmsas_20210317_1624/evt' is neither the name of a directory nor the name of a CIF
>export SAS_CCF='/blasco/users/gruppoXX/datadir/evt/ccf.cif'

The combination of RMF and ARF produces the input spectrum weighted by telescope area and detector efficiencies versus energy



14.GROUPING


In order to apply the χ^2 statistics (Gaussian distribution) you need to have at least 25 counts in each bin of your spectrum. Otherwise Cash statistics (Poisson distribution) is preferred (see also Statistics Tutorial).

```
grppha source_spectrum.fits pn_25.grp comm="chkey RESPFILE pn.rmf & chkey ANCRFILE pn.arf & chkey BACKFILE  
back_spectrum.fits & group min 25 & exit"
```

14.GROUPING

In order to apply the χ^2 statistics (Gaussian distribution) you need to have at least 25 counts in each bin of your spectrum. Otherwise Cash statistics (Poisson distribution) is preferred (see also Statistics Tutorial).

```
grppha source_spectrum.fits pn_25.grp comm="chkey RESPFILE pn.rmf & chkey ANCRFILE pn.arf & chkey BACKFILE  
back_spectrum.fits & group min 25 & exit"
```




see XSPEC tutorial...

15.LIGHT CURVE

<https://www.cosmos.esa.int/web/xmm-newton/sas-thread-timing>

A light curve is the plot of the flux of a source vs time.
It shows if and how the flux of the source varies during a certain time series.

The variability of a source can manifest on different time scales.



source+background lightcurve

pn

```
evselect table=pn_new.evt energycolumn=PI expression='#XMMEA_EP&&(PATTERN<=4)&& ((X,Y) IN circle(25910.5,25870.5,400)&& (PI in [2000:10000]))' withrateset=yes rateset="PN_source_lightcurve_raw.lc" timebinsize=100 maketimecolumn=yes makeratecolumn=yes
```

MOS 1

```
evselect table=mos1_new.evt energycolumn=PI expression='#XMMEA_EM&&(PATTERN<=12)&& ((X,Y) IN circle(25910.5,25870.5,400)&& (PI in [2000:10000]))' withrateset=yes rateset="MOS1_source_lightcurve_raw.lc" timebinsize=100 maketimecolumn=yes makeratecolumn=yes
```

MOS 2

The longer is the temporal
bin the lower is the
resolution but the higher
is the S/N.

```
evselect table=mos2_new.evt energycolumn=PI expression='#XMMEA_EM&&(PATTERN<=12)&& ((X,Y) IN circle(25910.5,25870.5,400)&& (PI in [2000:10000]))' withrateset=yes rateset="MOS2_source_lightcurve_raw.lc" timebinsize=100 maketimecolumn=yes makeratecolumn=yes
```

background lightcurve

pn

```
evselect table=pb_new.evt energycolumn=PI expression='#XMMEA_EP&&(PATTERN<=4)&& ((X,Y) IN circle(25910.5,25870.5,400)&&  
(PI in [2000:10000]))' withrateset=yes rateset="PN_light_curve_background_raw.lc" timebinsize=100 maketimecolumn=yes makeratecolumn=yes
```

MOS 1

```
evselect table=mos1_new.evt energycolumn=PI expression='#XMMEA_EM&&(PATTERN<=12)&& ((X,Y) IN circle(25910.5,25870.5,400)&&  
(PI in [2000:10000]))' withrateset=yes rateset="MOS1_light_curve_background_raw.lc" timebinsize=100 maketimecolumn=yes  
makeratecolumn=yes
```

The longer is the temporal
bin the lower is the
resolution but the higher
is the S/N.

MOS 2

```
evselect table=mos2_new.evt energycolumn=PI expression='#XMMEA_EM&&(PATTERN<=12)&& ((X,Y) IN circle(25910.5,25870.5,400)&&  
(PI in [2000:10000]))' withrateset=yes rateset="MOS2_light_curve_background_raw.lc" timebinsize=100 maketimecolumn=yes  
makeratecolumn=yes
```

source-background lightcurve: **epiclccorr**

pn

```
epiclccorr srctslist=PN_source_lightcurve_raw_lc eventlist=pn_new.evt outset=PN_lccorr_lc bkgtslist=PN_light_curve_background_raw_lc  
withbkgset=yes applyabsolutecorrections=yes
```


MOS 1

```
epiclccorr srctslist=MOS1_source_lightcurve_raw_lc eventlist=mos1_new.evt outset=MOS1_lccorr_lc  
bkgtslist=MOS1_light_curve_background_raw_lc withbkgset=yes applyabsolutecorrections=yes
```

MOS 2


```
epiclccorr srctslist=MOS2_source_lightcurve_raw_lc eventlist=mos2_new.evt outset=MOS2_lccorr_lc  
bkgtslist=MOS2_light_curve_background_raw_lc withbkgset=yes applyabsolutecorrections=yes
```

source-background lightcurve: `epiclccorr`




bin=100

The longer is the temporal
bin the lower is the
resolution but the higher
is the S/N.



bin=500

source-background lightcurve: epiclccorr



$$\chi^2_{\nu} = \frac{1}{\nu} \sum_{i=1}^n \frac{(c_i - \langle c \rangle)^2}{\sigma_i^2}$$

c_i observed counts in every temporal bin i;
 σ_i Poissonian error;
 $\langle c \rangle$ average count during the observation;
 $\nu=n-1$ degrees of freedom;

```
PLT> mo con  
1 CO: VAL( 1.000 ), SIG( 0.000 ), PL0( 0.000 ), PHI( 0.000 )?  
  
PLT> fit  
Fitting group 2, from 1.688E+03 to 9.038E+03  
Fitting 71 points in a band of 71.  
1.00000000  
( -3) W-VAR= 85.07  
( -4) W-VAR= 85.05  
0.178321302
```

Compute the probability of the result being due to chance

<https://www.fourmilab.ch/rpkp/experiments/analysis/chiCalc.html>

Calculate probability from X^2 and d

One of the most common chi-square calculations is determining, given the measured X^2 value for a set of experiments with a degree of freedom d , the probability of the result being due to chance. Enter the X^2 and d values in the boxes below, press the **Calculate** button, and the probability will appear in the Q box.

Given $X^2 = \boxed{85.05}$ and $d = \boxed{70}$ (71-1)

The chance probability, Q , is:

$1 - 0.1063 = 0.8937$ the source is variable at ~89% (the acceptance threshold for variability is 99.9%)

Summary

- **Download** the data from the archive
- **Data re-processing**
- **Data reduction:**
 - filtering for high particle background
 - selection of good time intervals (GTI)
 - selection of source and background extraction regions
- Check for the presence of **pileup**
- **Extraction** of source and background **spectra**
- Generate **RMF & ARF**
- **Grouping** (if possible...)
- (Background corrected) **light curve extraction**

