

On the $^{17}\text{O}/^{18}\text{O}$ ratio of post-AGB sources: canonical and non-canonical populations

Friday 14 June 2024 10:20 (20 minutes)

Stellar evolution and nucleosynthesis models are fundamental tools to derive parameters of stars from elemental atomic and isotopic abundance ratios. For evolved stars, the C/O/N elemental ratios and the $^{12}\text{C}/^{13}\text{C}$ and $^{16}\text{O}/^{17}\text{O}/^{18}\text{O}$ isotopic ratios provide very significant constraints to the otherwise elusive initial mass of the stars. In the case of post-AGB sources, the chemical composition of their expanding circumstellar envelopes can be derived from sensitive observations of rotational lines of several molecular species in the millimetre and sub-millimetre radio domains. The initial mass of post-AGB sources is a fundamental parameter to establish correlations with the main properties of their post-AGB envelopes (mass, momentum, kinetic energy, shape) and progress in our knowledge of their formation and evolution. In these sources, the C/O ratio can be easily constrained from the detection of C-bearing and O-bearing molecules other than CO, while the $^{17}\text{O}/^{18}\text{O}$ ratio is also straightforwardly determined from the relative strength of the optically thin rotational lines of C^{17}O and C^{18}O . However, the results obtained up to date are far from being clear.

In this presentation, we will review the status of the question, including new accurate $^{17}\text{O}/^{18}\text{O}$ ratio measurements for 13 targets, totalling 25 studied post-AGB envelopes: 15 O-rich sources (including the eight water fountains presented by Khouri *et al.* 2021) and 10 C-rich ones. Comparing the $^{17}\text{O}/^{18}\text{O}$ ratios and the C-rich/O-rich chemical composition with models for sources that have completed the AGB evolution, we find that for about 50% of the cases, observational data align with model predictions: these canonical sources include 50% of both O-rich and C-rich sources in the sample. As for the non-canonical sources, the O-rich ones, which present $^{17}\text{O}/^{18}\text{O}$ ratios above those expected for $\text{C}/\text{O} < 1$, can be explained by a premature interruption of their AGB evolution as a consequence of a quasi-explosive ejection of a large fraction of the initial mass. This hypothesis agrees with the suggestion that these envelopes form in the merging of common-envelope (or similar) events. The non-canonical C-rich sources, on the contrary, display $^{17}\text{O}/^{18}\text{O}$ ratios below the predictions for $\text{C}/\text{O} > 1$. We discuss possible explanations for this enigmatic behaviour, including the possibility that these sources are extrinsic C-rich stars, whose high C abundances result from previous mass accretion from a former higher mass C-rich AGB donor companion.

Primary author: ALCOLEA, Javier (Observatorio Astronómico Nacional (IGN/CNIG, Spain))

Co-authors: MASA ANDRÉS, Elisa (Observatorio Astronómico Nacional (OAN-IGN, Spain)); KHOURI, Theo (Chalmers University of Technology); SANTANDER-GARCÍA, Miguel (Observatorio Astronómico Nacional (IGN/CNIG, Spain)); Dr GALLARDO CAVA, Iván (Observatorio Astronómico Nacional (IGN/CNIG, Spain)); OLOFSSON, Hans (Space, Earth and Environment, Chalmers Univ. of Tech., Sweden); SANCHEZ CONTRERAS, Carmen (Centro de Astrobiología (CSIC-INTA, Spain)); BUJARRABAL, Valentín (Observatorio Astronómico Nacional (IGN/CNIG, Spain)); VLEMMINGS, Wouter (Chalmers University of Technology); Dr TAFOYA, Daniel (Chalmers University of Technology)

Presenter: ALCOLEA, Javier (Observatorio Astronómico Nacional (IGN/CNIG, Spain))

Session Classification: AGB and beyond: learning from the advanced phases