



Contribution ID: 65

Type: **Poster**

## **Kink-and-Disconnection Failed Eruption in 3D**

We report the first stereoscopic observations of HXR emission sources registered by STIX onboard Solar Orbiter and HXI on ASO-S. This is a case study of a two-stage failed eruption. First, it was slowed down due to a helical kink. However, the legs of the kinked structure started to reconnect and the second stage of eruption started. This eruption failed a few minutes later due to reconnection below the magnetic flux rope and confinement by overlying magnetic fields. We identified three X-ray sources located in the corona which are related to the reconnection sites and the magnetic cloud confined by overlying fields. Combining stereoscopic X-ray observations from STIX and HXI (31.5 degrees vantage point separation) with Differential Emission Measure (DEM) maps based on SDO/AIA observations we were able to locate X-ray source sites in the corona in the 3D space. The unveiled real geometry allowed us to estimate de-projected values of velocity and acceleration/deceleration which are extremely high. Moreover, real locations of HXR sources are not fully consistent with a standard solar flare scenario.

**Primary author:** MROZEK, Tomasz (Space Research Centre Polish Academy of Sciences)

**Co-authors:** Dr LI, Zhentong (Purple Mountain Observatory, Chinese Academy of Sciences); Prof. KARLICKÝ, Marian (Astronomical Institute ASCR); Dr CHRYSAPHI, Nicolina (Sorbonne Université, École Polytechnique, Institut Polytechnique de Paris, CNRS, Laboratoire de Physique des Plasmas (LPP)); Prof. SU, Yang (Purple Mountain Observatory, Chinese Academy of Sciences); Dr CHEN, Wei (Purple Mountain Observatory, Chinese Academy of Sciences); Prof. GAN, Weiqun (Purple Mountain Observatory, Chinese Academy of Sciences)

**Session Classification:** Coffee break and poster session 2

**Track Classification:** Multi-scale energy release, flares and coronal mass ejections