Contribution ID: 80 Contribution code: GAL

Type: Oral

## Investigating the hadron nature of high-energy photons with PeVatrons

Wednesday 4 September 2024 17:15 (15 minutes)

In high energy Gamma-Ray Astronomy with shower arrays the most discriminating signature of the photoninduced showers against hadron-induced cosmic-ray ones is the content of muons in the observed events. In the electromagnetic  $\gamma$ -showers the muon production is due to the dominant channels: photo-production of pions followed by the decay  $\pi \to \mu\nu$ , prompt leptonic decay of charmed particles in the shower, and electromagnetic pair production  $\gamma \to \mu^+ \mu^-$ .

The number of muons is typically a few percent of that in a hadron showers where muons are abundantly generated by charged pions decay.

In high energy photo-production process the photon exhibits an internal structure which is very similar to that of hadrons, with a small relative probability of order  $\alpha (\simeq 1/137)$ .

Indeed, photon-hadron interactions can be understood if the physical photon is viewed as a superposition of a bare photon and an accompanying small hadronic component which feels conventional hadronic interactions.

Information on photo-production  $\gamma p$  and  $\gamma \gamma$  cross-sections are limited to  $\sqrt{s} \leq 200$  GeV from data collected at HERA. Starting from  $E_{lab} \approx 100$  TeV the difference between different extrapolations of the cross sections increases to more than 50\% at  $E_{lab} \approx 10^{19}$  eV, with important impact in the observables used to select the photon-initiated air showers.

Recently, the LHAASO experiment opened the PeV-sky to observations detecting a number of PeVatrons in a background-free regime starting from about  $E_{lab} \approx 100$  TeV. This result provides a beam of pure high energy primary photons allowing to measure for the first time the photo-production cross section even at energies not explored yet.

The future air shower array SWGO in the Southern Hemisphere, where the existence of Super-Pevatrons emitting photons well above the PeV is expected, could extend the study of the hadron nature of the photons in the PeV region.

Primary author: Dr DI SCIASCIO, Giuseppe (INFN - Roma Tor Vergata)

Presenter: Dr DI SCIASCIO, Giuseppe (INFN - Roma Tor Vergata)

Session Classification: Parallel 1