

# StarDance: the non-canonical evolution of stars in clusters

*Wednesday 5 June 2024 09:25 (20 minutes)*

Thanks to their ubiquity, brightness, and the fact that they are made of stars with similar properties, star clusters have been used as astrophysical laboratories or test particles in an impressive range of research domains. However, we still do not understand fundamental details of their formation and evolution: a list of unsolved problems and apparently isolated mysteries has been accumulating over time, some standing since decades. Among them, the existence of multiple stellar populations in globular clusters, with different chemistry, has challenged generations of researchers. Given the mounting problems faced by the most favored scenarios to explain multiple populations, it is now time to revisit the foundations of our current thinking. New results show that: (i) the peculiarities in the chemistry of multiple populations are not limited to the oldest globular clusters; (ii) they can be transient in the evolution of individual cluster stars; and most importantly (iii) binary interactions and fast stellar rotation cannot be neglected in the study of star clusters and do have the capability to produce the observed chemistry. With StarDance, a new hypothesis will be tested: that multiple stellar populations and five other non-canonical stellar populations (extreme horizontal branch stars and hot sub-dwarfs; extended main sequence turn-offs; red stragglers and sub-subgiants; lithium-rich stars; and blue stragglers) are caused by the interplay between stellar rotation and binary interactions, that are greatly enhanced in the special environment of star cluster, with spectacular results.

## sessioni congresso

Stelle, popolazioni stellari e mezzo interstellare

**Primary author:** PANCINO, Elena (Istituto Nazionale di Astrofisica (INAF))

**Presenter:** PANCINO, Elena (Istituto Nazionale di Astrofisica (INAF))

**Session Classification:** Stelle, Popolazioni Stellari e Mezzo Interstellare