

# Neutron capture and total cross-section measurements on $^{94,95,96}\text{Mo}$ at n\_TOF and GELINA

Wednesday 11 June 2025 09:50 (20 minutes)

Cross-sections for neutron-induced interactions with molybdenum, in particular the neutron capture cross sections, play a significant role in various fields ranging from nuclear astrophysics to nuclear power plants and the development of innovative nuclear technologies. Molybdenum is found in pre-solar silicon carbide (SiC) grains and an accurate knowledge of its neutron capture cross section has a crucial role in stellar nucleosynthesis models, in particular in Asymptotic Giant Branch (AGB) stars. From the work of Liu et al., a deviation on the model predictions has been observed when using Mo cross section data from the two main KADoNiS versions, with KADoNiS 1.0 providing the better agreement with the grains data. In addition to its astrophysical role, molybdenum isotopes can be found as a fission product in fission power plants and the use of this material is under study for future improved reactors. This shows the importance of an accurate knowledge of the total and capture cross-section for molybdenum isotopes.

Experimental data in the literature for the capture cross-section of Mo isotopes suffer from large uncertainties. This is also reflected in the large uncertainties of the cross-sections recommended in the ENDF/B-VIII.0 library. Below 1 eV the relative uncertainty of the capture cross-section is above 18% for  $^{94}\text{Mo}$  and around 40% for  $^{96}\text{Mo}$ , while above 2 keV the uncertainties are in the order of 10-20% for  $^{94,95,96}\text{Mo}$ . The uncertainty on the capture cross section data in the libraires is also reflected in the uncertainty of the MACS (Maxwellian Averaged Cross Section) found in the latest version of KADoNiS. The presents uncertainties are on the level of 10% in the MACS at 30 keV for all the molybdenum isotopes, while an uncertainty below 5% would be needed to disentangle the observed discrepancies. One of the reasons for these large uncertainties is related to the absence of transmission data for enriched samples.

In this contribution the preliminary results of transmission and radiative capture measurements performed at n\_TOF (CERN, Switzerland) and GELINA (EC-JRC Geel, Belgium) will be presented. Moreover, the updated values of the MACS for  $^{94,95,96}\text{Mo}$  will be shown. The effect of these new preliminary values of the cross section in stellar nucleosynthesis calculations for AGB stars will be presented.

**Author:** MUCCIOLA, Riccardo (INFN Bari)

**Co-authors:** CRISTALLO, Sergio (Istituto Nazionale di Astrofisica (INAF)); KOPECKY, Stefan (European Commission, Joint Research Centre (JRC)); MANNA, Alice (CERN); MASSIMI, Cristian (INFN Bologna); MENGONI, Alberto (ENEA Bologna); MOENS, Andre (European Commission, Joint Research Centre (JRC)); PARADELA, Carlos (European Commission, Joint Research Centre (JRC)); SCHILLEBEECKX, Peter (European Commission, Joint Research Centre (JRC)); SIBBENS, Goedele (European Commission, Joint Research Centre (JRC)); VESCOVI, Diego (Istituto Nazionale di Astrofisica (INAF)); COLLABORATION, n\_TOF

**Presenter:** MUCCIOLA, Riccardo (INFN Bari)