




INAF
ISTITUTO NAZIONALE
DI ASTROFISICA

Chasing eruptive mass loss prior to superluminous supernovae

Anamaria Gkini


Supervisor: Ragnhild Lunnan

Co-supervisor: Jesper Sollerman




**Stockholm
University**

Massive stars will collapse and explode as CCSNe




Stars lose significant mass during their life




Stars can lose mass through various mechanisms


Stellar winds




Binary interaction




Eruptive mass-loss



CCSNe come in different flavours




CCSNe come in different flavours




What are superluminous supernovae?

- Transient events **10 — 100** times more luminous than normal CCSNe
- Rare events
 - 0.001% of CCSNe
- SLSNe-I are brighter than ~ -20 mag
- Their rise times range between **10 — 125 days**




Credits: Nicholl + 2015


Spectra can give insight into the SLSNe




Typical UV spectrum of SLSN-I




Typical UV spectrum of SLSN-I



Motivation





Credits: Lunnan +2018



Credits: Schulze +2024

Fast-moving CSM shell around SLSNe-I



Credits: Lunnan +2018

Fast-moving CSM shell around SLSNe


We collected X-shooter spectra of SLSNe

We observed 19 SLSNe with $z > 0.1$

Dedicated analysis of one objects in
Gkini+2024




Analysis of the sample in Gkini+ (in prep)




Two more SLSNe-I with Mg II signature

SN 2020xga




Credits: Gkini+2025



SN 2022xgc




Credits: Gkini+2025



We can extract CSM properties from the absorption lines




CSM shells expelled less than a year before CC




Credits: Gkini+2025

Expelled ~ 11 months before CC




Expelled ~ 5 months before CC

SN2018ibb the ejecta interacted with the CSM




Credits: Schulze+2024




Credits: Gkini+2025

Expelled ~ 9 months before CC

The Mg II system differs




Different line profiles



Various CSM properties

Eruptive mass loss is required to form CSM at this distances





Credits: Gkini+2025



Pulsational Pair Instability?

- The estimated ejection time put a limit on the He-core progenitor mass of $> 46\text{-}50 M_\odot$
- Geometry could perhaps give a hint of the expelling mechanism


Not all SLSNe show this signature




Not all SLSNe show this signature





Searching for eruptive mass loss...





Our sample



Is optical depth the key to detectability?



Two more objects !!!



Takeaways




Stockholm
University



- SN2020xga and SN2022xgc are SLSNe-I with Mg II absorption lines which indicate CSM shell around the SNe
 - The shells were expelled < 1 year before the explosion
- Pulsational Pair Instability of $> 50 M_{\odot}$ of He-core progenitors?
- **Six** objects have shown this feature. Rare events?
- Ongoing analysis could give insight into the mass loss before the explosion...

...AND THIS IS A
POSTDOC



© THE
UPTURNED
MICROSCOPE

