

SUPER-EDDINGTON ACCRETION UNDER THE MAGNIFIER: THE CASE OF ULTRALUMINOUS X-RAY SOURCES

FABIO PINTORE


(fabio.pintore@inaf.it, INAF/IASF Palermo)



Collaborators: C. Pinto, G. Rodriguez-Castillo, G. Israel, N. Pinciroli Vago, D. Walton, F. Fuerst, S.Motta, A. D'Aì, M. Del Santo, R. Salvaterra, E. Ambrosi, A. Wolter, L. Zampieri, Wladimiro Leone, Luciano Burderi and many more

ULTRALUMINOUS X-RAY SOURCES

- Point-like X-ray sources
- Off-Nuclear (i.e. no supermassive BH)
- $L_x >> 10^{39}$ erg/s $>> L_{edd}$ of 10 M_{Sun} BH
- X-ray Luminosities as high as 10⁴¹⁻⁴² erg/s



ULTRALUMINOUS X-RAY SOURCES

OBSERVATIONS OF THE X-RAY SOURCES IN THE NEARBY Sc GALAXY M33

KNOX S. LONG

Columbia Astrophysics Laboratory, Columbia University

SANDRO D'ODORICO European Southern Observatory

PHILIP A. CHARLES¹


Space Sciences Laboratory, University of California

AND

MICHAEL A. DOPITA

Mt. Stromlo and Siding Spring Observatory, Australian National University Received 1980 December 29; accepted 1981 February 25

ABSTRACT




THE ASTROPHYSICAL JOURNAL, 325:544-562, 1988 February 15 © 1988. The American Astronomical Society. All rights reserved. Printed in U.S.A.

THE X-RAY EMISSION OF M81 AND ITS NUCLEUS

G. FABBIANO

Harvard-Smithsonian Center for Astrophysics Received 1987 April 14; accepted 1987 July 29





overy of ybroeck. r source. s greater ral arms, also find oution of A coms a more an those 1 or also sities. h energy model is ir source pret this s for the $N_{\rm H}$ is far he explactrum, if :-10 keV

More than ~2000 ULXs have been catalogued (e.g. Walton et al. 2022)!

However, mostly are poorly known in terms of spectral and temporal properties


ULXs are thought to be accreting compact objects in binary system






ULTRALUMINOUS X-RAY SOURCES

ULXs were initially considered as a rescaled version of the Galactic BH binaries, and therefore accreting sub-eddington.




To sustain luminosities higher than 10³⁹ erg/s up to 10⁴¹⁻⁴² erg/s, the BH mass should be increased up to very large masses!

M ~ 100-10⁵ Msun




HLX-1 in ESO 243-49 is the best candidate to host an IMBH (Farrel et al. 2009, Webb et al. 2013)




 $L_x -> 10^{42} \text{ erg/s}$

 $M_{BH} >> 10^3 M_{sun}$





ATCA observed a variable radio emission at 5 GHz and 9 Ghz, likely associated to a jet




Spectral evolution compatible with a hard-to-soft

A jet was observed in radio in the source M31 ULX-1 during its 2013 outburst






Clear spectral evolution resembling the hard-to-soft transition of Galactic BH binaries



Swift/XRT long-term lightcurve of the only registered outburst of this ULX.


A CRACK IN THE IMBH SCENARIO




ULXs

Hard Ultraluminous
Soft Ultraluminous
Broadened disc

Fractional variability (F_{var}) significantly higher in ULX soft ultraluminous states, unlike Galactic BH binaries







Hysteresis cycle?


ULXs do not follow it...


ULX spectral properties are show a pletora of behaviours


(e.g. Gurpide et al. 2021)












ULTRALUMINOUS X-RAY SOURCES




Pintore et al. (2021)

Pintore et al. (2021)

O.5 1.0 2.0 5.0 Energy (keV)

Transitions from soft ultraluminous to hard ultraluminous states observed in several ULXs, but different global spectral behaviour




Observed Mass Ranges of Compact Objects Stellar **Intermediate Mass Supermassive Black Hole Black Hole Black Hole Neutron** Star White Super-Eddington accretion! Dwarf a committee de la committe del la committe de la co 10 100 1,000 10,000 100,000 1,000,000 **Object Mass** (Relative to the Sun) Colbert & Mutshotzky (1999), Madau & Rees (2004)


Portgies Swartz et al. (2004)

- Pulsation have been discovered in at least 6 extragalactic ULXs using observations with high counting statistics (XMM-Newton and NuSTAR)
- P_{spin} in the range ~ 0.4 20 s


M82 X-2 (BACHETTI ET AL. 2014) Period (ms) -03 - 10 keV 03 - 30 keV 10 - 30 keV 56680 56710 56690 56700 56720 MJD





NGC 7793 P13 (ISRAEL ET AL.2017)



- Pulsation have been discovered in at least 6 extragalactic ULXs using observations with high counting statistics (XMM-Newton and NuSTAR)
- P_{spin} in the range ~ 0.4 20 s
- PULXs orbital properties suggest periods of a few days and giant OB companion stars








SUPER-CRITICAL ACCRETION REGIME

To explain luminosities well above 10³⁹ erg/s, super-Eddington accretion has to be taken into account;

At super-Eddington accretion, a standard disc is not expected;

Radiatively emitted outflows with an equatorial geometry

 $R_{sph} \sim \dot{M}_0$




See Poutanen et al. (2007)

MHD:


Ohsuga et al. (2009)
Kawashima et al. (2014)
Romavoa et al.(2014)
Mushtukov et al. (2015)
Mushtukov et al. (2016)
Kawashima et al. (2016)


SUPER-CRITICAL ACCRETION REGIME





Soft lags, i.e. soft emission lags the hard one, observed in several ULXs!

- Lags of hundreds of seconds (compatible with those observed in AGN)
- Possible reprocessing of hard photons from the funnel of wind/outflows.













SUPER-CRITICAL ACCRETION REGIME









Optical/IR/radio nebulae (>100 pc in size).
Inflated by outflows?






Belfiore, Esposito, FP et al. (2019)



A nebula was also found in X-rays around the pulsating ULX named NGC 5907 ULX-1, in one Chandra observation


ORBITAL PROPERTIES IN PULXS




Observed a secular orbital evolution in M82 X-2 which supports the existence of outflows in this source (Bachetti et al. 2022).

MAGNETIC FIELDS IN PULXS

High magnetic field (B > 10¹⁴ G) can reduce the scattering cross-section and allows for super-Eddington accretion over the NS polar caps. $\frac{\sigma_{\rm B1}}{\sigma_{\rm T}} \approx \sin^2\theta + \left(\frac{E_{\gamma}}{E_{\rm cyc}}\right)^2 \cos^2\theta$






 $rac{\sigma_{
m B2}}{\sigma_{
m T}} pprox \left(rac{E_{\gamma}}{E_{
m cyc}}
ight)^2, {
m for} \; rac{E_{\gamma}}{E_{
m cyc}} \ll 1,$

High B field (B > 10^{14} G)

Low B field (B < 10¹³ G)

MAGNETIC FIELDS IN PULXS

High magnetic field (B > 10^{14} G) can reduce the scattering cross-section and allows for super-Eddington accretion over the NS polar caps.



Pieces of evidence support lower dipolar B-fields (< 10¹³ G) in ULXs:

- presence of outflows;
- PULXs should be continuously in propeller;
- sinusoidal shape of the pulsed emission

Quadrupolar magnetic field components?

ULTRALUMINOUS X-RAY SOURCES: OPEN QUESTIONS


- What's the nature of the companion stars (HMXBvs LMXB)?
- How are the outflows properties?
- How do outflows imprint on the surrounding medium?
- What's the compact object mass in ULXs?
- What's the BH vs NS ratio in the ULX population?
- ▶ Unknown NS magnetic field (dipolar vs quadrupolar)? Magnetar-like (10^{14-15} G) or more standard (10^{11-13} G)?
- Can the magnetic field truncate the disc before the spherization radius? Outflows are inhibited in PULXs? Not really, implying non-magnetar like magnetic fields.
- Is there any geometrical beaming of the emission?
- ▶ How does the spin evolve with time (spin-up/spin-down)?
- Is the large flux variability seen in PULXs due to propeller?

ULTRALUMINOUS X-RAY SOURCES: OPEN QUESTIONS

- What's the nature of the companion stars (HMXBvs LMXB)?
- How are the outflows properties?
- How do outflows imprint on the surrounding medium?
- What's the compact object mass in ULXs?
- What's the BH vs NS ratio in the ULX population?
- ▶ Unknown NS magnetic field (dipolar vs quadrupolar)? Magnetar-like (10^{14-15} G) or more standard (10^{11-13} G)?
- Can the magnetic field truncate the disc before the spherization radius? Outflows are inhibited in PULXs? Not really, implying non-magnetar like magnetic fields.
- Is there any geometrical beaming of the emission?
- ▶ How does the spin evolve with time (spin-up/spin-down)?
- Is the large flux variability seen in PULXs due to propeller?

- Only 6 PULXs over the total population of ~2000 ULXs
- ▶However, PULXs are ~20-30% of the ULX population which has high quality observations to detect pulsations (~30 sources)
- ▶ Are the vast majority of ULXs hosting BH then?
- Increase the sample of PULXs. However, pulsations are intermittent in PULXs! Search for PULX-like sources amongst ULXs

NEUTRON STARS VS BLACK HOLES IN THE ULX POPULATION



Cyclotron line at $4.5 \text{ keV} (3.8\sigma \text{ detection})$ in a Chandra spectrum of M51 ULX-8


 $(B/10^{12} G) = E/(11.6(1 + z)^{-1}) -> for electrons$

 $(B/10^{15} G) = E/(6.3(1 + z)^{-1}) -> for protons$


Estimated magnetic field of $B < 10^{12} \, G$ (Brightman et al. 2018; Middleton, Brightman, Pintore et al. 2019)

No pulsation detected yet...


- Pulsating ULXs:
 - All transient sources (with a possible by-modality)





- Pulsating ULXs:
 - All transient sources (with a possible by-modality)
 - Spectral shapes



- Pulsating ULXs:
 - All transient sources (with a possible by-modality)
 - Spectral shapes
 - Super-orbital modulations of tens-tohundreds of days








NGC 7793 P13 (Fuerst et al.2018)


HUNT FOR PULX CANDIDATES

- Pulsating ULXs:
 - All transient sources (with a possible by-modality)
 - Spectral shapes
 - Super-orbital modulations of tens-tohundreds of days
 - (Quasi-periodic) flaring activity




M51 ULX-7 (Imbrogno et al. 2024)


FUTURE DEVELOPMENTS: NEWATHENA



NewAthena will help for both high resolution and broadband spectroscopy in the ULX field



- The vast majority of ULXs is expected to be stellar mass compact objects accreting super-Eddington -> an extreme version of Galactic X-ray binaries
- Still many things to unveil, in particular how super-Eddington accretion processes work in pulsating ULXs
- Outflows are a marked footprint of ULXs and almost ubiquitous in these sources
- Winds -> responsible for most of the temporal and spectral properties observed in ULXs
- Outflows feedbacks on the galactic environment
- Investigating super-Eddington accretion in ULXs can help to understand the process of growing for supermassive BHs in AGN
- Needs for population studies. How many BH are in the ULX population?



Thanks for the attention