

Tracing Cosmic Magnetism with Gamma-ray Bursts

Davide Miceli
INFN Padova

davide.miceli@pd.infn.it

IFPU

11/02/2025

About me

- Born in Modena, 15 October 1993
- PhD at University of Udine (but living in Trieste 😊)



Multi-wavelength afterglow numerical code and data analysis of MAGIC very high energy emission from gamma-ray bursts (GRBs)

- A little bit travel around for postdocs/positions:
 - Local operator specialist for MAGIC and LST telescopes, living in La Palma, Canary Island
 - Postdoctoral fellowships:
 - LAPP, Annecy, France
 - University of Padova, Italy
 - Non-permanent PNNR position at Istituto Nazionale di Fisica Nucleare (INFN), Padova, **CTA+ project**

About me


IGMF: when did everything start?

- 2021 University of Padova 'Probing cosmic magnetism with high-energy astrophysics'
→ investigate the capability of Cherenkov telescopes (MAGIC, CTAO) of studying IGMF from GRBs detected at Very High Energy (VHE, $E > 100$ GeV)
- Exploring a refined approach to assess the impact of intrinsic GRB properties (energetics, geometry, distance) in IGMF studies
- Collaborators: E. Prandini, P. Da Vela, L. Nava, G. Ghirlanda, ..?

Background and expertise:


- **Gamma-ray Cherenkov telescopes** (MAGIC, LST, CTAO): data analysis; simulations; maintenance activities; follow-up transient sources; currently Coordinator of the MAGIC Transient Working Group
- **Gamma-ray Bursts**: observations in the gamma-ray GeV-TeV domain; theoretical interpretation and modeling of multi-wavelength (radio to gamma-rays) afterglow emission;
- **Intergalactic magnetic field studies in gamma-ray domain with GRBs**

A (non-exhaustive) map of topics involved in this talk





Gamma-ray Bursts


- Transients (ms – days/weeks)
- Cosmological ($z \sim 2$)
- Bright ($L_{\text{iso}} 10^{49} - 10^{53} \text{ erg s}^{-1}$)




Gamma-ray Bursts



Gamma-ray Bursts: Prompt phase





Gamma-ray Bursts: afterglow phase



Is there emission in the VHE domain ($E > 100$ GeV) in GRBs?

HE emission (< 100 GeV) properties:

- Almost consistent with synchrotron radiation (synchrotron burnoff limit)
- No spectral cut-off identified (shock microphysics uncertainties, non-uniform magnetic fields)




Observations from keV to TeV domain

keV → GeV → TeV



Credit: ESA



Credit: NASA



Credit: NASA

Space telescopes



Credit: Robert Wagner/MAGIC Collaboration




Ground-based observatories

Credit: LHAASO

Observations of GRBs: TeV domain

Long story short:


- We need to know **where** to look (space telescopes trigger and send alert with position uncertainty of arcmins)
- We need to be **fast** (follow-up starting at order of min or even sec)




Observations of GRBs: TeV domain

- **Imaging Atmospheric Cherenkov Telescopes (IACTs) [30-50 GeV – 10s TeV]**
 - Narrow field of view (FoV) ($3-5^\circ$) \rightarrow need external alert + automatic alert system + fast repointing
 - Small duty cycle (10%)
 - Low energy threshold: 30-50 GeV
 - Better sensitivity than EAS up to a few TeV
- **Shower front arrays [100s GeV – >100s TeV]:**
 - Wide field of view \rightarrow no repointing needed
 - Large duty cycle (100%)
 - High energy threshold: 100s GeV – TeV \rightarrow extragalactic background light (EBL) absorption
 - Large collection area \rightarrow can reach 100s TeV and above

Imaging Atmospheric Cherenkov Telescopes (IACTs)



Credit: Robert Wagner/MAGIC Collaboration



Credit: Klepser, DESY, H.E.S.S. collaboration



Credit: VERITAS



Credit: Tomohiro Inada, CTAO



Imaging Atmospheric Cherenkov Telescopes (IACTs)




Hunting GRBs in the VHE domain: a non-homogeneous story

From Berti & Carosi 2022

- 1989: first detection of VHE source (Crab Nebula) by IACT (Whipple telescope)
- 1991-2000 BATSE, EGRET operations
- 1994-95: first follow-ups by IACT (Whipple, 9 GRBs) → delays from $T_0 \sim 2-56$ minutes
- 1996: launch of Beppo-Sax
- 1994-00: several EAS searches for TeV/PeV emission from GRBs (including hint of Milagrito on 970417A)
- 2004 launch of Swift → fundamental for alerts and successfully rapid repointing
- 2004: MAGIC-I, H.E.S.S. phase-I (4x12m) and ARGO-YBJ started operations and GRB follow-up
- 2007: VERITAS started operations and GRB follow-up
- 2007-08 AGILE (07) and Fermi (08) were launched
- 2009: MAGIC-II started operations → big sensitivity improvement
- 2012 HESS phase-II: add 28 m telescope → transient dedicated telescope


Decades of searches for the VHE emission




Berti et al., ICRC 2021

Decades of searches for the VHE emission

Cherenkov telescope observations: **only upper limits until 2019**



Aleksic et al. 2014



Aliu et al., 2014

GRB 190114C: 1^o announcement from IACTs

TITLE: GCN CIRCULAR
NUMBER: 23701
SUBJECT: MAGIC detects the GRB 190114C in the TeV energy domain
DATE: 19/01/15 01:56:36 GMT
FROM: Razmik Mirzoyan at MPI/MAGIC <Razmik.Mirzoyan@mpp.mpg.de>

R. Mirzoyan (MPP Munich), K. Noda (ICRR University of Tokyo),
E. Moretti (IFAE Barcelona), A. Berti (University and INFN Torino),
C. Nigro (DESY Zeuthen), J. Hoang (UCM Madrid), S. Micanovic
(University of Rijeka), M. Takahashi (ICRR University of Tokyo),
Y. Chai (MPP Munich), A. Moralejo (IFAE Barcelona) and the MAGIC
Collaboration report:

On January 14, 2019, the MAGIC telescopes located at the Observatorio Roque de los Muchachos on the Canary island of La Palma, detected very-high-energy gamma-ray emission from GRB 190114C (Gropp et al.,

GRBs at VHE: the current status

5 GRBs detected at $> 5\sigma$ (afterglow phase)

	T_{90} s	$E_{\gamma,iso}$ erg	z	T_{delay} s	E_{range} TeV	IACT (sign.)
160821B	0.48	1.2×10^{49}	0.162	24	0.5-5	MAGIC (3.1σ)
180720B	48.9	6.0×10^{53}	0.654	3.64×10^4	0.1-0.44	H.E.S.S. (5.3σ)
190114C	362	2.5×10^{53}	0.424	57	0.3-1	MAGIC ($> 50\sigma$)
190829A	58.2	2.0×10^{50}	0.079	1.55×10^4	0.18-3.3	H.E.S.S. (21.7σ)
201015A	9.78	1.1×10^{50}	0.42	33	0.14	MAGIC (3.5σ)
201216C	48	4.7×10^{53}	1.1	56	0.1	MAGIC (6.0σ)
221009A	289	1.0×10^{55}	0.151	0-3000	0.3-13	LHAASO (250σ)


GRBs at VHE: the current status

2 GRBs detected at $\sim 3\sigma$ (afterglow phase)


	T_{90} s	$E_{\gamma,iso}$ erg	z	T_{delay} s	E_{range} TeV	IACT (sign.)
160821B	0.48	1.2×10^{49}	0.162	24	0.5-5	MAGIC (3.1σ)
180720B	48.9	6.0×10^{53}	0.654	3.64×10^4	0.1-0.44	H.E.S.S. (5.3σ)
190114C	362	2.5×10^{53}	0.424	57	0.3-1	MAGIC ($> 50\sigma$)
190829A	58.2	2.0×10^{50}	0.079	1.55×10^4	0.18-3.3	H.E.S.S. (21.7σ)
201015A	9.78	1.1×10^{50}	0.42	33	0.14	MAGIC (3.5σ)
201216C	48	4.7×10^{53}	1.1	56	0.1	MAGIC (6.0σ)
221009A	289	1.0×10^{55}	0.151	0-3000	0.3-13	LHAASO (250σ)

Population of GRBs in VHE domain


- **Broadband intrinsic properties:**
 - span more than 3 orders of magnitude in $E_{\gamma,iso}$
 - Span 2 orders of magnitude in terms of L_{VHE}
 - ranging in redshift between 0.079–1.1
- **X-ray – TeV connection:**
 - similar fluxes and decay slopes
 - similar amount of radiated power (L_{VHE} 15-60% L_{X-ray})
- **Data modeling:**
 - SSC suggested (not conclusive)
 - no preferences on constant/wind-like medium
 - $\epsilon_e \sim 0.1$, $\epsilon_B \sim 10^{-5}-10^{-3}$, $\xi < 1$



VHE responsible radiation mechanism in GRBs




HESS Coll., 2021




Salafia et al., 2021

Population of GRBs in VHE domain: the role of redshift

long GRBs



short GRBs




adapted from Nava, 2021

Open question for VHE in GRBs


- Is TeV emission a universal component in GRBs?
- Is SSC the responsible radiation mechanism or is there an interplay among radiation mechanisms? (synchrotron radiation, SSC, photo-hadronic interactions)
- How does the VHE spectrum evolve with time?
- Prompt VHE emission?
- Several open field in GRB physics: shock microphysics, environmental conditions at the burst site, acceleration process efficiency, orientation-dependency and connection with GW

How GRBs impact on IGMF studies?

Razzaque et al. (2004)
Ichiki et al. (2008)
Takahashi et al. (2008)
Murase et al. (2009)



An extended and time-delayed component due to IGMF deflection + CMB reprocessing



How GRBs impact on IGMF studies?

Intrinsic source properties


Gamma-ray propagation (EBL, pair scatter, IC) and magnetic field models

Instrument sensitivity


GRB intrinsic source spectrum (spectral and temporal evolution) → **input for gamma-ray propagation**

How can gamma-ray probe IGMF properties (B strength and correlation length λ_B)?


Extended emission



Time-delayed "pair-echo" emission





SED signatures



Pair-echo after end of TeV afterglow emission

Plaga 1995
Murase et al. 2008



GeV source emission

+

pair echo cascade emission with non-negligible IGMF

Interesting for AGN flares or transient sources

Search for the time-delayed 'pair-echo' cascade emission

$$E_{rep} \sim 0.32 \left(\frac{E_\gamma}{20 \text{ TeV}} \right)^2 \text{ TeV}$$

$$F_{delay} \sim F_0 \frac{T}{T_{delay} + T}$$

$$T_{delay} \propto B^2 E_\gamma^{-5/2}$$


$$T_{delay} \propto B^2 E_\gamma^{-2} \lambda_B$$

$$\lambda_B \gg \lambda_{IC}$$

$$\lambda_B \ll \lambda_{IC}$$

Neronov et al. 2009

Batista et al. 2021



The advantage of using GRBs for IGMF studies?

- GRBs timescales of emission (hours, days in gamma-ray domain) → reduced pollution from primary source
- GRB duty cycle → relaxed assumption
- GRB intrinsic brightness and cosmological nature → increase redshift horizon
- GRBs discovered at TeV energies → IACTs can play a role (GRB intrinsic spectrum + pair-echo)




An independent verification of IGMF studies with AGNs

An opportunity to explore pair-echo signal without source contamination at different energy ranges

Pair-echo emission from TeV-detected GRBs with IACTs

GRB190114C ($z = 0.42$)




Spectral energy distribution

- Primary GRB emission
- Secondary emission
- Observational time: 3 hours starting from 2400 s after trigger burst
- MAGIC and CTAO sensitivity derived and rescaled in time ($S \propto (1/vt)$)

Pair-echo emission from TeV-detected GRBs with IACTs


GRB190114C ($z = 0.42$)

Scaled GRB190114C ($z = 0.2$)



Pair-echo emission from TeV-detected GRBs with IACTs

GRB221009A ($z = 0.151$)





DM, Da Vela, Prandini, 2024

- Extend observations for at least 3 hours after GRB detection
- GRBs observations can probe IGMF strengths in the $10^{-17} - 10^{-19} \text{ G}$ \rightarrow competitive with most stringent AGN results


Pair-echo emission from TeV-detected GRBs with IACTs

GRB221009A ($z = 0.151$)

Scaled GRB221009A ($z = 1.0$)




Pair-echo emission + GRB afterglow



- Afterglow emission can vary of several order of magnitudes
- X-ray afterglow displays unclear features (plateaus, flares, steep decay, jet breaks)
- How **GRB intrinsic properties** impact IGMF studies? (is it true that brightest and nearby GRBs are best choices?)

Pair-echo emission + GRB afterglow convolution


Extrapolate GRB properties (spectrum and time evolution) from a simulated GRB




On going study with P. Da Vela, G. Ghirlanda, L. Nava

Pair-echo emission + GRB afterglow convolution

We estimated the pair-echo LCs from a simulated GRB




Pair-echo emission + GRB afterglow convolution



- Assuming a smaller redshift ($z=0.15$) with same GRB properties
- Add a “jet break” at 10⁴ s (light curve steepening of a factor $\propto t^{-1}$)

Pair-echo emission + GRB afterglow convolution



- Pair-echo emission becomes competitive with GRB afterglow at late times for $B > 10^{-19}$ G

Currently exploring the **impact of GRB properties** (energies, distance, geometry) on a sample of simulated GRBs

Next generation: Cherenkov Telescope Array Observatory (CTAO)

CTAO North (Alpha configuration)



4 LSTs (23 m)
9 MSTs (12 m)

CTAO South (Alpha configuration + CTA+)



2 LSTs
14 MSTs
37+5 SSTs



Next generation: Cherenkov Telescope Array Observatory (CTAO)



Next generation: Cherenkov Telescope Array Observatory (CTAO)

CTAO North status May 2024 →
commissioning in 2025/2026




Credit: M.Mariotti

Next generation: Cherenkov Telescope Array Observatory (CTAO)

CTAO upgrades:


- a lower **energy threshold** (<30 GeV)
- a larger **effective area** at multi-GeV energies ($\sim 10^4$ times larger than Fermi-LAT at 30 GeV)
- a **rapid slewing** capability (180 degrees azimuthal rotation in 20 s).
- a **full sky coverage**
- a **broader energy range**: 20 GeV- 300 TeV



CTAO for GRBs: an upcoming revolution

CTAO revolution:

- x10 sensitivity to gamma-ray signal
- x10⁴ sensitivity to short-term (< 10⁴ s) emission below 250 GeV with respect to Fermi-LAT
- Intermediate arrays at both sites operative by 2027; full array by 2031-2032
- GRB detection rates (Inoue et al. 2013): > 1-4 GRBs/year/site (but to be updated with new VHE GRB discovery!)



Conclusions (+ future prospects)

- Discoveries in the past 5 years have open a new observational window for GRBs: the **TeV domain**
- Results have shown that (to some extent) VHE emission is compatible with **broadband intrinsic properties** of GRBs (energetics, luminosity, distance) and **broad observational requirements** (timing, VHE range) → universality of TeV component?
- So far modeling reproduced the VHE component in the context of the **SSC emission** from external forward shock scenario, but still not conclusive results (Synchrotron, EIC, p-synchrotron)
- **GRBs** can provide new **independent and complementary verification of IGMF studies** in gamma-ray domain, circumventing limitations and assumptions of AGN studies
- Impact of **GRBs intrinsic source features** still unclear (distance, brightness, intrinsic spectrum shape and features, time evolution, jet break) → currently under study!
- Current and future generation instruments (**CTAO, SWGO**) are ready in the game, improvements are under investigation, 15 years of continuous improvements lead to first GRB VHE detections



Finanziato
dall'Unione europea
NextGenerationEU



Ministero
dell'Università
e della Ricerca




Italiadomani
PIANO NAZIONALE
DI RIPRESA E RESILIENZA




BACKUP SLIDES



Shower front arrays





Next generation: the Southern Wide-field Gamma-ray Observatory (SWGGO)



Barres de Almeida, 2022

Gamma-ray bursts with SWGO



Search for the time-delayed 'pair-echo' cascade emission

$$E_{rep} \sim 0.32 \left(\frac{E_\gamma}{20 \text{ TeV}} \right)^2 \text{ TeV}$$

$$F_{\text{delay}} \sim F_0 \frac{T}{T_{\text{delay}} + T}$$

$$T_{\text{delay}} \propto B^2 E_\gamma^{-5/2} \quad \lambda_B \gg \lambda_{IC}$$
$$T_{\text{delay}} \propto B^2 E_\gamma^{-2} \lambda_B \quad \lambda_B \ll \lambda_{IC}$$

- 100s GeV photons experience shorter delays (\sim hrs/days) than GeV photons (\sim weeks/yrs)
- Weak B field ($10^{-17} - 10^{-21}$ G) are compatible short delays
- Stronger B are compatible with longer delays (and a more diluted cascade)

Neronov et al. 2009

Batista et al. 2021

Search for the time-delayed 'pair-echo' cascade emission


$$E_{rep} \sim 0.32 \left(\frac{E_\gamma}{20 \text{ TeV}} \right)^2 \text{ TeV}$$

$$F_{delay} \sim F_0 \frac{T}{T_{delay} + T}$$

$$T_{delay} \propto B^2 E_\gamma^{-5/2} \quad \lambda_B \gg \lambda_{IC}$$

$$T_{delay} \propto B^2 E_\gamma^{-2} \lambda_B \quad \lambda_B \ll \lambda_{IC}$$

Neronov et al. 2009
Batista et al. 2021



Search for the time-delayed 'pair-echo' cascade emission


$$E_{rep} \sim 0.32 \left(\frac{E_\gamma}{20 \text{ TeV}} \right)^2 \text{ TeV}$$

$$F_{delay} \sim F_0 \frac{T}{T_{delay} + T}$$

$$T_{delay} \propto B^2 E_\gamma^{-5/2} \quad \lambda_B \gg \lambda_{IC}$$

$$T_{delay} \propto B^2 E_\gamma^{-2} \lambda_B \quad \lambda_B \ll \lambda_{IC}$$

Neronov et al. 2009
Batista et al. 2021




GRB 190114C -- Timeline

- $E_{\gamma,iso} \sim 2.5 \times 10^{53}$ erg
- $z = 0.42$

MAGIC detection info:


- $T_{delay} \sim 57$ s
- $> 50\sigma$ in 20 minutes
- detection up to 40 min
- 0.3 - 1 TeV energy range
- moon conditions and $Zd > 50$




GRB 190114C – Light curve

VHE light curve:

- No evidences for breaks, cut-offs or irregular variability → afterglow emission
- Similar decay and radiated power in soft X-ray – GeV and TeV domain



GRB 190114C – VHE SED



GRB 190114C -- interpretation

Acceleration timescale (Bohm level)


$$t'_L = \frac{r_L}{c} = \frac{\gamma m_e c}{eB'}$$

e-Synchrotron cooling timescale

$$t'_{\text{cool}} = \frac{E'}{|dE'/dt'|} = \frac{6\pi m_e c}{\gamma \sigma_T B'^2}$$



Synchrotron burnoff limit

$$\epsilon_{\text{sync,max}} \sim 100 \left(\frac{\Gamma_b}{1000} \right) \text{ GeV}$$



MAGIC Coll. et al., 2019

GRB 190114C -- interpretation



GRB 190114C -- modeling

Synchrotron + Synchrotron Self-Compton (SSC) from external forward shock

Observed
 No γ - γ opacity
 EBL-deabsorbed

MAGIC soft spectrum:

- Klein-Nishina
- γ - γ internal absorption

GRB afterglow parameters:


$$E_k \gtrsim 3 \times 10^{53} \text{ erg}$$

$$\epsilon_e \sim 0.05 - 0.15$$

$$\epsilon_b \sim 0.05 - 1 \times 10^{-3}$$

$$n \sim 0.5 - 5 \text{ cm}^{-3}$$

$$p \sim 2.4 - 2.6$$




GRB 190829A


- $E_{\gamma,iso} \sim 2.0 \times 10^{50}$ erg
- $z = 0.079$

H.E.S.S. detection info:


- $T_{obs} \sim 4.3 - 55.9$ hrs
- $21.7\sigma, 5.5\sigma, 2.4\sigma,$
- $0.18 - 3.3$ TeV energy range



GRB 190829A



HESS Coll., 2021




Salafia et al., 2021

GRB 201216C

- $E_{y,iso} \sim 4.7 \times 10^{53}$ erg
- $z = 1.1$

MAGIC detection info:


- $T_{delay} \sim 56$ s
- 6σ in 20 minutes
- 0.07 – 0.2 TeV energy range



GRB 201216C

Parameter	Range	Best fit value
E_k [erg]	$10^{50} - 10^{54}$	4×10^{53}
θ_{jet} [degrees]	0.5 - 3	1
Γ_0	80-300	180
n_0 [cm^{-3}] ($s = 0$)	$10^{-2} - 10^2$	-
A_\star ($s = 2$)	$10^{-2} - 10^2$	2.5×10^{-2}
p	2.05 - 2.6	2.1
ϵ_e	0.01-0.9	0.08
ϵ_B	$10^{-7} - 10^{-1}$	2.5×10^{-3}

Strong indication in favour of a wind-like medium




GRB 221009A

- $E_{\gamma,iso} 1 \times 10^{55}$ erg
- $z = 0.15$



LHAASO detection info:

- $> 250\sigma$ in 230 – 3000 s
- 0.3 – 13 TeV energy range



LHAASO Coll. et al., 2023

GRB 221009A




GRB 180720B

- Long GRB
- $E_{\gamma,iso} \sim 6.0 \times 10^{53}$ erg
- $z = 0.654$

H.E.S.S. detection info:

- Tdelay ~ 10 hrs
- $> 5.3\sigma$ in 2 hrs
- 0.1 – 0.44 TeV energy range




GRB 160821B

- short GRB
- $E_{\gamma,iso} \sim 1.2 \times 10^{49}$ erg
- $z = 0.162$

MAGIC info:

- Tdelay ~ 24 s
- 3σ in 4 hrs
- 0.5 - 5 TeV energy range
- moon conditions, dedicated analysis





GRB 190114C

Time bin [seconds after T_0]	Energy flux [erg cm ⁻² s ⁻¹]	Spectral index
62 – 100	$[5.64 \pm 0.90 \text{ (stat)}^{+3.24}_{-3.22} \text{ (sys)}] \cdot 10^{-8}$	$-1.86^{+0.36}_{-0.40} \text{ (stat)}^{+0.12}_{-0.21} \text{ (sys)}$
100 – 140	$[3.31 \pm 0.67 \text{ (stat)}^{+2.71}_{-1.84} \text{ (sys)}] \cdot 10^{-8}$	$-2.15^{+0.43}_{-0.48} \text{ (stat)}^{+0.25}_{-0.32} \text{ (sys)}$
140 – 210	$[1.89 \pm 0.36 \text{ (stat)}^{+1.72}_{-0.94} \text{ (sys)}] \cdot 10^{-8}$	$-2.31^{+0.47}_{-0.54} \text{ (stat)}^{+0.15}_{-0.22} \text{ (sys)}$
210 – 361.5	$[7.54 \pm 1.60 \text{ (stat)}^{+6.46}_{-4.41} \text{ (sys)}] \cdot 10^{-9}$	$-2.53^{+0.53}_{-0.62} \text{ (stat)}^{+0.22}_{-0.24} \text{ (sys)}$
361.5 – 800	$[3.10 \pm 0.70 \text{ (stat)}^{+1.20}_{-2.36} \text{ (sys)}] \cdot 10^{-9}$	$-2.41^{+0.51}_{-0.65} \text{ (stat)}^{+0.27}_{-0.34} \text{ (sys)}$
800 – 2454	$[4.54 \pm 2.04 \text{ (stat)}^{+7.66}_{-1.96} \text{ (sys)}] \cdot 10^{-10}$	$-3.10^{+0.87}_{-1.25} \text{ (stat)}^{+0.75}_{-0.24} \text{ (sys)}$
62 – 2454 (time integrated)	-	$-2.22^{+0.23}_{-0.25} \text{ (stat)}^{+0.21}_{-0.26} \text{ (sys)}$



GRB 190114C

Time bin [seconds after T_0]	D11	F08	FI10	G12
62 – 100	$-1.86^{+0.36}_{-0.40}$	$-2.04^{+0.36}_{-0.40}$	$-1.81^{+0.36}_{-0.40}$	$-1.95^{+0.36}_{-0.39}$
100 – 140	$-2.15^{+0.43}_{-0.48}$	$-2.32^{+0.43}_{-0.48}$	$-2.09^{+0.43}_{-0.48}$	$-2.23^{+0.42}_{-0.48}$
140 – 210	$-2.31^{+0.47}_{-0.54}$	$-2.48^{+0.47}_{-0.54}$	$-2.25^{+0.47}_{-0.54}$	$-2.39^{+0.47}_{-0.53}$
210 – 361.5	$-2.53^{+0.53}_{-0.62}$	$-2.69^{+0.52}_{-0.61}$	$-2.46^{+0.52}_{-0.61}$	$-2.60^{+0.52}_{-0.61}$
361.5 – 800	$-2.41^{+0.51}_{-0.65}$	$-2.58^{+0.51}_{-0.64}$	$-2.34^{+0.51}_{-0.64}$	$-2.49^{+0.51}_{-0.64}$
800 – 2454	$-3.10^{+0.87}_{-1.25}$	$-3.20^{+0.83}_{-1.20}$	$-2.96^{+0.83}_{-1.20}$	$-3.08^{+0.82}_{-1.19}$
62 – 2454 (time integrated)	$-2.22^{+0.23}_{-0.25}$	$-2.39^{+0.23}_{-0.25}$	$-2.15^{+0.23}_{-0.25}$	$-2.29^{+0.23}_{-0.24}$



GRB 190114C

Event	redshift	T_{delay} (s)	Zenith angle (deg)
GRB 061217	0.83	786.0	59.9
GRB 100816A	0.80	1439.0	26.0
GRB 160821B	0.16	24.0	34.0
GRB 190114C	0.42	58.0	55.8

Population of GRBs at VHE: what we thought vs what we discovered

“Mandatory” requirements:

- low zenith angles (energy threshold below ~ 100 GeV)
- dark nights
- small delays from T_0
- low z
- highly energetic events

GRB190114C: zenith $>55^\circ$, Moon conditions

GRB160821B: Moon conditions


GRB180720B, GRB190829A: $T_{\text{delay}} \sim \text{hrs/days}$

GRB201216C: $z = 1.1$

GRB190829A, GRB201015A, GRB160821B: $E_{\text{y,iso}} \sim 10^{49} - 10^{50}$ erg

Population of GRBs in VHE domain

- Broadband intrinsic properties:
 - span more than 3 orders of magnitude in $E_{\gamma,iso}$
 - Span 2 orders of magnitude in terms of L_{VHE}
 - ranging in redshift between 0.079–1.1



DM & Nava, 2022, adapted from D'Avanzo et al. 2014

Open question: degeneracy of afterglow parameters

	E_k erg	ϵ_e	ϵ_B	n cm^{-3}	p	ζ_e	θ_j rad
Hess Coll. (SSC)	2.0×10^{50}	0.91	$5.9-7.7 \times 10^{-2}$	1.	2.06-2.15	1.	/
Hess Coll. (Sync)	2.0×10^{50}	0.03-0.08	≈ 1	1.	2.1	1.	/
Salafia + 2021	$1.2-4.4 \times 10^{53}$	0.01-0.06	$1.2-6.0 \times 10^{-5}$	0.12-0.58	2.01	$< 6.5 \times 10^{-2}$	0.25-0.29
Zhang + 2021	9.8×10^{51}	0.39	8.7×10^{-5}	0.09	2.1	0.34	0.1

	E_k erg	ϵ_e	ϵ_B	n cm^{-3}	p	ζ_e
MAGIC Coll.	$\gtrsim 3 \times 10^{53}$	0.05-0.15	$0.05-1 \times 10^{-3}$	0.5-5	2.4-2.6	1
Wang + 2019	6×10^{53}	0.07	4×10^{-5}	0.3	2.5	1
Asano + 2020	10^{54}	0.06	9×10^{-4}	1	2.3	0.3
Asano + 2020	10^{54}	0.08	1.2×10^{-3}	0.1 (wind)	2.35	0.3
Joshi + 2021	4×10^{54}	0.03	0.012	2×10^{-2} (wind)	2.2	1
Derishev + 2021	3×10^{53}	0.1	$2-6 \times 10^{-3}$	2	2.5	1

	E_k erg	$\log(\epsilon_e)$	$\log(\epsilon_B)$	$\log(n)$ cm^{-3}	p	ζ_e	θ_j rad
MAGIC Coll.	$10^{51}-10^{52}$	[-1; -0.1]	[-5.5; -0.8]	[-4.85; -0.24]	2.2-2.35	1	/
Troja + 2019	$10^{50}-10^{51}$	[-0.39; -0.05]	[-3.1; -1.1]	[-4.2; -1.7]	2.26-2.39	1	0.08-0.50
Zhang + 2021 (SSC)	3×10^{51}	-0.52	-5	-1.3	2.3	0.5	0.15
Zhang + 2021 (EIC)	2×10^{51}	-0.3	-6	-1	2.5	0.1	0.1

A future challenge for VHE: X-ray Flares


Signatures of X-ray flares can be found in the GeV-TeV domain?

Wang et al. 2006


He et al. 2012

Wang et al. 2013

1/3 of GRBs display X-ray flare episodes (Chincarini et al. 2010, Margutti et al. 2011)




DM, Nava 2022



Adapted from Fioretti et al. 2019 <https://www.ing.unipi.it/it/azioni/73> Ricerca

Population of GRBs in VHE domain: the role of redshift



Dominguez et al., 2011
(similar for other EBL models)

$z \lesssim 0.1 - 0.2$

- F_{att} relevant above 300 GeV
- $F_{\text{att}} \sim 90\%$ at 1 TeV

$z = 0.4$


- $F_{\text{att}} \sim 50\%$ at 0.2 TeV
- $F_{\text{att}} \sim 99.5\%$ at 1 TeV

$z = 1.1$


- $F_{\text{att}} \sim 95\%$ at 0.2 TeV

Theoretical expectations from GRBs in the VHE domain

Synchrotron self-Compton (SSC) emission has been predicted for GRB afterglows: nature candidate for VHE domain (Meszaros et al. 1994; Zhang et al. 2001; Sari et al. 2001; Meszaros et al. 2004; Fan et al. 2008; Galli et al. 2008; Nakar et al. 2009; Xue et al. 2009; Piran et al. 2010)




Sari, Esin, 2001




Afterglow modeling: external forward shock scenario

Decelerating blastwave interacting with the circumburst external medium



Afterglow modeling: external forward shock scenario

Jet dynamics




See Blandford & Mckee, 1976; Nava et al., 2014


Afterglow modeling: external forward shock scenario

Relativistic shocks in GRB afterglow

See
Sari et al.
1998,
Panaitescu
et al. 2000
Granot et al.
2002




Afterglow modeling: external forward shock scenario



See
Sari et al.
1998,
Panaitescu
et al. 2000
Granot et al.
2002

Intergalactic Magnetic field (IGMF) studies

IGMF studies investigate a 2D parameter space: Correlation Length (λ_B) – IGMF Strength (B)



Results on IGMF are typically given considering two regimes:

- Long correlation length ($\lambda_B \gg \lambda_{IC}$)
(motion in homogeneous B, ballistic e^\pm)
- Short correlation length ($\lambda_B \ll \lambda_{IC}$)
(diffusion in angle, diffusive e^\pm)


Pair-echo emission + GRB afterglow convolution

$$F_c(E, t) = \int_0^\infty \int_E^\infty G(E_0, E, t - \tau, \tau) F_s(E_0, t - \tau) dE_0 d\tau$$

Cascade Flux

Kernel describing the distribution in energy and time of the cascade signal

"Variability pattern" (Source intrinsic properties and time evolution)



Gamma-rays for IGMF studies: Methods

How gamma-ray can probe IGMF properties (B strength and correlation length λ_B)?

- Method I : search for extended emission

- A “smoking gun” for IGMF discovery
- Size and shape depend on IGMF strength and source parameters (jet opening and orientation)

$$\Theta_{\text{ext}} \propto B E_{\gamma}^{-1} \quad \lambda_B \gg \lambda_{\text{IC}}$$

$$\Theta_{\text{ext}} \propto B E_{\gamma}^{-3/4} \lambda_B^{1/2} \quad \lambda_B \ll \lambda_{\text{IC}}$$

