


# Euclid Legacy Science: Milky Way and Local Universe

Francesca Annibali<sup>1</sup> (MWRSP deputy lead)  
and  
Leslie Hunt<sup>2</sup> (LU co-lead),  
on behalf of the MWRSP and LU working groups

<sup>1</sup> INAF-OAS  
<sup>2</sup> INAF-OAA

# Tracing galaxy evolution with resolved stars



Color-magnitude diagrams are powerful tools to age-date stars





# Tracing galaxy evolution with resolved stars


Color-magnitude diagrams are powerful tools to age-date stars





The deeper the CMD,  
the higher the  $z$  we can probe

# The Euclid revolution: high angular resolution & large FoV

**LBT (8.4 m)**  
seeing limited



**HST (2.4 m)**  
diffraction limited




Resolving individual stars to deep magnitudes  
requires **high angular resolution** in crowded fields!




# The Euclid revolution high angular resolution

**LBT** (8.4 m)  
seeing limited




**HST** (2.4 m)  
diffraction limited



But many many pointings  
required with HST to cover  
what Euclid can do in just  
one shot!

Resolving individual stars to deep magnitudes  
requires **high angular resolution** in crowded fields!

# Euclid Wide Survey will image many nearby systems




Several Galactic GCs

Not the bulk of the GC system toward the Galactic Center




# Euclid Wide Survey will image many nearby systems



Huge number of  
MW dwarf  
satellites:  
dSphs and UFDs

Not M31+M33 &  
Satellites


# Euclid Wide Survey will image many nearby systems



Many late  
type dwarfs in  
the outskirts of  
the LG and just  
beyond



# Euclid Wide Survey will image many nearby systems



M81 group,  
M101 Group,  
Sculptor group

Not Cen A, Not  
Virgo

# The Milky Way and Resolved Stellar Populations (MWRSP) SWG

**Leads:** B. Goldman, E. Balbinot, **F. Annibali** (Deputy-lead)

## Work Packages:

Italian  
contribution

**WP-Photometry:** Stellar populations age, Z, distances  
(**F. Annibali**/ M. Rejkuba)

**WP-Spectroscopy:** star classification and radial velocities (M. Manteiga/ G. Torralba Elípe)

**WP-Galactic Globular clusters:** extended halos of GCs  
(**D. Massari**)

**WP-Extra -Galactic Globular clusters:** Joint WP with the Local Universe SWG (Ariane Lançon)

**WP-Ultra-Cool dwarfs:** (**R. L. Smart** / C. Reyle)





# The Milky Way and Resolved Stellar Populations (MWRSP) SWG

**Leads:** B. Goldman, E. Balbinot, **F. Annibali** (Deputy-lead)

## Approved DR1 Key Projects :

Italian  
contribution

DR1-KP-MWRSP1: Globular Clusters

Coordinators : **E. Dalessandro**, S. Larsen

DR1-KP-MWRSP2: Resolved stellar peripheries

Coordinators: A. Ferguson, E. Balbinot

DR1-KP-MWRSP3: The Magellanic Clouds

Coordinators: M. Romero




Euclid: ERO – AGGIORNARE)





# The Periphery of Milky Way Globular Clusters

D. Massari, E. Dalessandro, et al.(2024)


- ERO program targeting two MW GCs, NGC 6397 and NGC 6254 (PI Massari)
- Imaging in  $I_E, Y_E, J_E, H_E$ , depth similar to Wide Survey
- Resolving stellar peripheries of GCs, looking for tidal tails induced by interaction with the MW



Euclid: ERO – Globular Clusters-image processing by J.-C. Cuillandre





Density map from resolved star counts provides evidence for tidally induced morphological distortions in NGC 6255



# High precision Astrometry and Photometry with Euclid

M. Libralato, L. Bedin, M. Griggio, D. Massari, et al. (2024)

- Test on ERO data of NGC 6397 (PI Massari) with specific pipeline for high precision astrometry and photometry



- Astrometric precision of **0.7 mas** in Euclid VIS and **3 mas** in NISP
- Comparison of Euclid astrometry with Gaia DR3 (epoch 2016) provides higher precision PMs than Gaia!


Potentially, multi-epoch Euclid astrometry would allow for high precision PMs down to  $I_E \sim 26$ !!



# Resolved Stellar Populations in Nearby Galaxies

L. Hunt, F. Annibali, et al.(2024)


Euclid ERO Nearby Galaxy Showcase Program (PI Hunt)




- Selected from WISE Extended Source Catalog
- Angular size < Euclid FoV (to fit in 1 ROS)
- Ancillary data (e.g. HI)

# The extended stellar component in the dwarf IC10

F. Annibali, A. Ferguson, P. Sanchez-Alarcon, P. Dimauro, et al, in prep.




Spatial distribution of RGB stars







# The extended stellar component in the dwarf IC10

F. Annibali, A. Ferguson, P. Sanchez-Alarcon, P. Dimauro, et al, in prep.




Spatial distribution of RGB stars




Stellar component much more extended than previously thought!

Stellar profile requires two distinct components





# The extended stellar component in IC10

F. Annibali, A. Ferguson, P. Sanchez




Spatial distribution of RGB stars



Stellar component much more extended than previously thought!

Stellar profile requires two distinct components





# The Local Universe(LU) SWG


**Leads:** M. Fabricius, **L. Hunt**, F. Marleau, T. Saifollahi (Junior Deputy)

Two main science themes for the LU motivated by Euclid's capabilities:

- (1) **low surface brightness features** because of their importance for probing remnants of hierarchical structural assembly in galaxies (stellar streams, tidal debris, intracluster light, dwarf galaxies, ...)
- (2) superb spatial resolution enables measuring **physical properties** on parsec /10s of pc scales, and characterization of **extragalactic globular clusters (EGCs)**



Euclid will image to unprecedented depths faint, low surface brightness galaxies, stellar streams/tidal debris, and globular clusters in galaxies up to  $z \sim 0.2$ , enabling new galaxy demographics and new distance calibrations through surface-brightness fluctuations



Euclid: ERO – Dwarf galaxies in the Perseus galaxy cluster: EC, Marleau+ (2025)

# Approved pre-launch Key Projects and Q1/DR1 Key Projects

LU Key Project I: Opening Euclid's window on low surface brightness

Coordinators: **Enrica Iodice**, Reynier Peletier, **Leslie Hunt**

LU Key Project II: Simulating the Euclid View of the Local Universe

Coordinators: Johan Knapen, **Crescenzo Tortora**

KP-LU-DR1-1: Euclid's low surface brightness view of the nearby universe from DR1

Coordinators: **Leslie Hunt**, Francine Marleau

KP-LU-DR1-2: Stellar populations in nearby galaxies from DR1

Coordinators: **Daniela Carollo**, **Crescenzo Tortora**, Karina Voggel

KP-LU-DR1-3: The Fornax Cluster and Supercluster: New insights from Euclid

Coordinators: Pierre-Alain Duc, **Enrica Iodice**

KP-LU-DR1-4: Surface Brightness Fluctuation distances with Euclid

Coordinators: **Rebecca Habas**, Simona Mei

Italian  
contribution



# Work Packages:

WP-PPZ: Physical parameters and photometric redshifts (Marco Scodeggio / Crescenzo Tortora)

WP-FAR: Distance estimates (Simona Mei)

WP-DET: Detection Algorithms (Paola Dimauro)

WP-MOR: Morphology measurements (Louis Quilley)

WP-LBG: Dwarf and low surface brightness galaxies (Pierre-Alain Duc / Reynier Peletier)

WP-DIF: Diffuse emission (Fernando Buitrago)

WP-EGC: Extragalactic globular clusters (Ariane Lançon)


Italian  
contribution

# Distance measurements with Surface Brightness Fluctuations

Lead: Michele Cantiello (INAF-OA Teramo )

Rebecca Habas (post-doc supported by INAF Large grant EDGE, PI Hunt)

See R. Habas POSTER!



MW Globular Cluster  
~10 kpc

Messier 32  
~770 kpc

Messier 49  
~16500 Mpc


- Preferred targets: elliptical galaxies with smooth SB profile
- Number of stars/pixel has Poisson fluctuations, **more distant galaxies appear smoother**
- Observed "mottling" is directly related to the target distance. Measurement error on this mottling ~0.05 mag or less

SBF def.: Ratio 2nd to 1st moments of stellar luminosity function

$$f_{SBF} \equiv \bar{f} \equiv \sum_i n_i f_i^2 / \sum_i n_i f_i$$

$f^2 \rightarrow$  bright stars weight more

$$\bar{m} \sim \langle m_{RGB} \rangle$$



Tonry & Schneider 1988

Tonry, Ajhar & Luppino 1990


Blakeslee, Ajhar & Tonry 1999..

Goal: to develop a self-contained, automated code for the community, adapted to *Euclid*. Conservatively, >50 Mpc with Euclid.




# Distance measurements with Surface Brightness Fluctuations


original image




residual image



fluctuations (2D covariance)



power  
spectrum of  
azimuthally  
averaged  
fluctuations



## Fornax cluster for the first tests:

- close enough to measure SBF
- several galaxies already have SBF measurements (e.g., Blakeslee et al. 2009)
- dwarf candidates already identified


R. Habas, M. Cantiello, et al. in prep.

## Measure SBF in $I_E$ band:

- better resolution
- Lanczos3 interpolation for stacking (interpolation method important)
- **SPOIL ALERT: bi-linear interpolation kills SBF signal**

See R. Habas POSTER!

## schematic power spectrum



# Physical Parameters and Redshift (PPZ) WP Abdurro'uf, Tortora et al. 2025

**Leads:** Crescenzo Tortora, Marco Scodeggio


## Objectives:

- photometric redshifts and integrated stellar populations for galaxies at  $z < 0.2$
- spatially-resolved stellar population maps for nearby galaxies
- Test performance of spatially resolved-SED fitting on mock datasamples (e.g. EURISKO, used in Humphrey, Cunha, Bisigello, Tortora, et al 2015)


### Results:

1. stellar mass density well recovered with Euclid-only bands
2. improved performance when adding ancillary data from other surveys

Euclid only



GALEX + LSST + Euclid




# PPZ WP: Plans for DR1

DR1 contains ~5000 local galaxies with distances.

## Objectives:

- Colour maps
- Application of the spatially-resolved SED fitting and machine learning tools
- Morphological characterization




*IYH colour images of Q1 galaxies with distances from HECATE catalog (Kovlakas et al. 2021) –  
Image credits: Tortora/Euclid*



# Euclid as a Low Surface Brightness Machine

Comparison of the **surface brightness limit** ( $3\sigma$ ,  $10 \times 10$  arcsec<sup>2</sup>) for the Euclid/VIS Wide Survey and Deep Survey, compared with a selection of deep optical and near-infrared surveys.

Although a few surveys may go deeper, they do so over very small areas. Euclid will probe the LSB universe on a huge area of the sky, opening up a new parameter space for LSB science




Taken from EC, Borlaff+ (2022), see also Scaramella+ (2022)

# Low Surface Brightness Data Labs pilot study

P. Dimauro et al., *in prep.*

EC pipeline optimized for the main Science goal of Euclid: Cosmology





Need an LSB friendly pipeline

# Low Surface Brightness Data Labs pilot study

P. Dimauro et al., *in prep.*

EC pipeline optimized for the main Science goal of Euclid: Cosmology



See Paola Dimauro  
hands-on session on  
Wed July 2<sup>nd</sup>!




# Low Surface Brightness Data Labs pilot study

P. Dimauro et al., *in prep.*

Now available LSB pipeline-reduced VIS and NISP images in Datalabs: [datalabs.esa.int](https://datalabs.esa.int)



Euclid team\_workspace  
Euclid-Consortium data  
LSB\_reduction  
VIS  
EDF-N  
EDF-S  
• EDFN\_Mosaic.table.fits  
• EDFS\_Mosaic.table.fits



See Paola Dimauro  
hands-on session on  
Wed July 2<sup>nd</sup>!

# Low Surface Brightness Data Labs pilot study

Comparison MER and DataLabs LSB pipelines: **ESO202-023 post merger** (courtesy J. Miro)




See Paola Dimauro  
hands-on session on  
Wed July 2<sup>nd</sup>!

MER pipeline tends to oversubtract background








Euclid VIS 1-sigma asinh limiting surface brightness (LSB science performance) and the nearby universe up to  $z=0.03$

- Wide Survey region of interest : 17 Kdeg.<sup>2</sup> compliant with a 15 Kdeg.<sup>2</sup> survey
- Deep Fields : North=10 deg.<sup>2</sup>, Fornax=10 deg.<sup>2</sup>, South=23 deg.<sup>2</sup> (+2 mag.)
- ⇒ metric reflecting the ultimate science LSB performance based on profile extraction



AGGIORNARE: Euclid will image to unprecedented depths faint, low surface brightness galaxies, stellar streams/tidal debris, and globular clusters in galaxies up to  $z \sim 0.2$ , enabling new galaxy demographics and new distance calibrations through surface-brightness fluctuations

