

Extra-galactic distances for the next generation surveys: the surface brightness fluctuation method



Osservatorio Astronomico d'Abruzzo

Gabriele Riccio

**With: Michele Cantiello (CL),
Rebecca Habas, Gabriella
Raimondo, Matteo Canzari**

ITA-INA-S6) Directable SW contribution for the Galaxies SC: Tools for the measurement of surface brightness fluctuations on LSST data

Coordinator



Michele Cantiello

Post-doc on LSST

3 years (ending March 2027)



Gabriele Riccio

Software Engineer



Matteo Canzari

Post-doc on Euclid

3 years (ending May 2027)

Rebecca Habas

Stellar population models


Gabriella Raimondo

Extra-Galactic distances: what's the tension?

- **Reliable distances** are required to understand most observed objects & phenomena
- **Errors in distances** imply significant fractional uncertainties in derived quantities

Masses (e.g., central black holes, dark matter halos); Physical sizes;
 Dynamical time scales; Star Formation Rates; Ages...

- **Precise extragalactic distances** play a key role in the H_0 controversy, a major issue in modern cosmology




Pop II ladder

Geometry → TRGB


TRGB → SBF

SBF → galaxy distances,
dwarf membership,
Hubble flow

Riess +2020



Surface Brightness Fluctuation (SBF)



MW Globular Cluster
~10 kpc

Messier 32
~770 kpc


Messier 49
~17 Mpc

Slide Basics: R. Kudritzki/M.Cantiello
Images: J. Blakeslee

SBF def.: Ratio 2nd to 1st moments of stellar luminosity function

$$f_{SBF} \equiv \bar{f} \equiv \frac{\sum_i n_i f_i^2}{\sum_i n_i f_i}$$

$f^2 \rightarrow$ bright stars weight more



Tonry & Schneider 1988
Tonry, Ajhar & Luppino 1990
Blakeslee, Ajhar & Tonry 1999...



Preferred targets: Elliptical galaxies with smooth surface brightness profiles.



Number of unresolved stars/pixel has Poisson fluctuations, more distant galaxies appear smoother.



Observed “mottling” is directly related to the target distance. Measurement error on this mottling 0.05 mag or less



No dust or non-stellar-count fluctuations!

SBF with LSST (promising)

<i>5σ-depth (mag)</i>	26.3-26.8 (i-band)
<i>FWHM (arcsec)</i>	0.6-0.7
<i>Filters</i>	ugirzy
<i>Sky coverage (sq. deg)</i>	~18,000
<i>D_{Max}</i>	~70 Mpc (for the best candidates)

Build the most accurate
3D map of the local
universe.

Measure H0 independently
from Cepheid and SNe Ia.

- Thousands of optimal/good candidates
- $\gg 10^3$ non-optimal candidates for which distances to 10-20% can be derived (in some cases no other distance indicator might be available)



Measuring SBF is a long process...



Many steps involved + **Heavy user interaction**

=


t ~ hours

Stellar counts fluctuations $P_f = P_0 - P_r$


$$\bar{m}_{band} = -2.5 \log(P_f) + m_{zp, band}$$




(Usually done manually)




P_r : Spurious fluctuation term



P_0 : Total fluctuation term




\bar{m}



FAST-SBF


(**F**lexible **A**utomated **S**elf-contained **T**ool for **SBF**)

- **Flexible**: the pipeline is optimise to work on the major large-field surveys
- **Automated**: the user interaction is reduced at the minimum (with the aim to remove it completely), maintaining reliable results.
- **Self-contained**: all the steps of the entire analysis are carried out by one tool.
- **Fast**: takes from few seconds to tens of minutes (depending on the source) to completely analyse a galaxy.




FAST-SBF: first results

Subaru HSC-SSP (LSST precursor)



Riccio+2025

Riccio+2025




Distance and comparison with SSP models



$$\bar{M} = \alpha \times color + \beta$$

β tied to **Cepheid** or **TRGB**.
Also validated through simple stellar population models.


First result on DP1 (ComCam)



DP2 (LSSTcam) expected by mid July 2026



Catalyst post-doc alliance



FAST-SBF in Teramo:
a workshop to measure distances with
Surface Brightness Fluctuations

Teramo, Italy
20 - 23 July 2026



SOC: John Blakeslee, Enzo Brocato, Michele Cantiello,
Matteo Canzari, Rebecca Habas, Joe Jensen,
Gabriella Raimondo, Gabriele Riccio



Summary

Science:






- Precise on single red/bright targets: ~5%
- $\Delta d/d \sim 15/20\%$ on the other extreme, dwarf galaxies
- Easy observations
- Well understood and modelled origin of the signal
- The method needs to be adapted to run in “production mode” to deliver robust results quickly and with minimal human intervention
- The method is promising for both distance indicator and stellar population diagnostic, especially for dwarfs




FAST-SBF


In-kind(2024-2026):

- Develop a tool for the SBF analysis capable of handling big data.
- Testing the pipeline on LSST precursors.
- Data preview 1
- Data preview 2 (July - September 2026)
- Data Release (>2027)



Regarding
SBF in the
context of
stellar
populations...










The TRGB-SBF project with the JWST

NIRCAM: ideal layout



Jensen+25

$$H_0 = 73.8 \pm 0.7 \pm 2.3 \text{ km/s/} \\ \text{Mpc}$$





SBF with LSST: maximum distance



- Main limiting factor in measuring SBF at large distances: 'spurious fluctuations', mostly from globular clusters host in the galaxy.



- At the present LSST survey specs: in final coadds SBF will be measured in all bands. Largest distances (~ 50 Mpc) reached in i, z and y bands.
- Lucky cases (good FWHM, bright red target, ...) possibly 70 Mpc.

Name	(g-i) (mag)	\bar{m}_i (mag)	$\bar{m} - \bar{M}$ (mag)	d (Mpc)
(1)	(2)	(3)	(4)	(5)
NGC 5813	1.206 (0.004)	31.58 (0.08)	31.92 (0.12)	24.2 (1.3)
PGC 053521	0.991 (0.007)	30.92 (0.08)	32.02 (0.12)	25.4 (1.4)
PGC 053587	0.998 (0.013)	30.98 (0.28)	32.07 (0.29)	26.0 (3.5)
PGC 053636	0.921 (0.017)	30.74 (0.10)	32.01 (0.13)	25.2 (1.6)
PGC 1165764	0.455 (0.022)	29.35 (0.09)	32.04 (0.13)	25.6 (1.5)
PGC 1192611	0.798 (0.026)	30.73 (0.22)	32.23 (0.24)	28.0 (3.2)
PGC 1193898	0.988 (0.019)	30.95 (0.09)	32.08 (0.13)	26.0 (1.5)
PGC 1205406	0.897 (0.039)	30.57 (0.18)	31.90 (0.20)	23.9 (2.3)
PGC 1208589	0.887 (0.028)	30.45 (0.17)	31.79 (0.20)	22.8 (2.1)
PGC 1230503	0.920 (0.016)	31.08 (0.11)	32.36 (0.14)	29.6 (1.9)
PGC 184851	1.050 (0.007)	31.09 (0.07)	32.02 (0.12)	25.4 (1.4)
PGC 3350761	1.105 (0.033)	31.68 (0.21)	32.46 (0.23)	31.1 (3.4)
PGC 3350757	1.047 (0.023)	31.18 (0.14)	32.16 (0.17)	27.0 (2.1)
PGC 3350778	0.932 (0.018)	30.72 (0.06)	31.98 (0.11)	24.9 (1.2)
PGC 4018458	0.872 (0.037)	31.05 (0.14)	32.42 (0.17)	30.5 (2.4)
PGC 4018466	0.936 (0.043)	30.96 (0.23)	32.20 (0.25)	27.6 (3.18)
PGC 4074773	0.858 (0.017)	31.04 (0.16)	32.25 (0.18)	28.1 (2.4)
NGC 5839	1.131 (0.004)	31.50 (0.17)	32.17 (0.19)	27.2 (2.4)
PGC 1199471	0.927 (0.023)	30.75 (0.08)	32.01 (0.12)	25.2 (1.4)
PGC 4009173	0.688 (0.028)	29.82 (0.16)	31.55 (0.18)	20.4 (1.7)
NGC 5831	1.133 (0.008)	31.48 (0.09)	32.17 (0.13)	27.1 (1.6)
PGC 1190315	1.043 (0.07)	30.79 (0.09)	31.78 (0.14)	22.7 (1.5)
PGC 1216386	0.918 (0.017)	30.89 (0.08)	32.17 (0.12)	27.1 (1.5)
NGC 4753	0.958 (0.002)	30.70 (0.07)	31.86 (0.11)	23.5 (1.2)
PGC 1150193	0.363 (0.049)	28.97 (0.14)	32.01 (0.17)	25.2 (2.1)
PGC 135809	0.667 (0.053)	29.63 (0.18)	31.37 (0.20)	18.8 (1.8)
PGC 183384	0.780 (0.013)	30.0 (0.10)	31.51 (0.13)	20.1 (1.2)
PGC 3295853	0.750 (0.041)	30.20 (0.23)	31.77 (0.25)	22.5 (2.6)
PGC 44066	0.5012 (0.013)	29.46 (0.19)	31.72 (0.21)	22.1 (2.1)
NGC 4690	0.975 (0.003)	31.515 (0.06)	32.63 (0.11)	33.5 (1.7)
PGC 1115603	0.736 (0.026)	30.67 (0.26)	32.26 (0.27)	28.4 (3.6)
PGC 1124896	0.570 (0.025)	30.22 (0.34)	32.22 (0.35)	27.8 (4.5)
IC 0745	0.828 (0.008)	30.21 (0.06)	31.63 (0.11)	21.17 (1.1)
GAMA559879	0.546 (0.037)	30.01 (0.056)	32.09 (0.11)	26.2 (1.3)
GAMA583929	0.586 (0.047)	30.78 (0.64)	32.73 (0.64)	35.1 (10.4)







Rubin Legacy Survey of Space and Time (LSST)

Telescope (constraint)	Band	Depth ^(a) (mag)	GCLF peak ^(b)	GCLF offset ^(c)	d_{Max} (Mpc)	Optimal/good candidates	$\delta H_0/H_0$ (%) optimal/good
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Rubin (Dec $\leq 15^\circ$)	<i>i</i>	26.8	-8.5	+0.5	70	50/1500	1.2/<0.5
Euclid ($ b \geq 23^\circ$)	H_{Euclid}	24.0	-8.6	-1.5	55	20/600	1.0/<0.5
Roman ^(e)	J_{F129}	28.0	-8.3	-1.0	230	1500/>5000	0.7/0.5 ^(g)
Roman ^(e)	H_{F158}	28.0	-8.6	-1.5	300	1600/>5000	0.7/0.5 ^(g)
JWST ^(e)	J_{F115W}	27.5	-8.3	-1.0	180	900/>5000	0.9/0.5 ^(g)
JWST ^(e)	H_{F150W}	27.7	-8.6	-1.5	290	1600/>5000	0.7/0.5 ^(g)
JWST ^(e)	K_{F200W}	27.8	-8.1	-2.0	310	1600/>5000	0.7/0.5 ^(g)
ELT ^(f) (Dec $\leq 0^\circ$)	<i>K</i>	28.5	-8.1	-2.0	400	750/>4000	1.0/0.7 ^(g)

	LSST
5σ -depth (mag)	26.3-26.8 (i-band)
FWHM (arcsec)	0.6-0.7
Filters	<i>ugrizy</i>
Sky coverage (sq. deg)	~20,000



Euclid



Telescope (constraint)	Band	Depth ^(a) (mag)	GCLF peak ^(b)	GCLF offset ^(c)	d_{Max} (Mpc)	Optimal/good candidates	$\delta H_0/H_0$ (%) optimal/good
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Rubin (Dec < 15°)	<i>i</i>	26.8	-8.5	+0.5	70	50/1500	1.2/<0.5
Euclid ($b \geq 23^\circ$)	H_{Euclid}	24.0	-8.6	-1.5	55	20/600	1.0/<0.5
Roman ^(e)	J_{F129}	28.0	-8.3	-1.0	230	1500/>5000	0.7/0.5 ^(g)
Roman ^(e)	H_{F158}	28.0	-8.6	-1.5	300	1600/>5000	0.7/0.5 ^(g)
JWST ^(e)	J_{F115W}	27.5	-8.3	-1.0	180	900/>5000	0.9/0.5 ^(g)
JWST ^(e)	H_{F150W}	27.7	-8.6	-1.5	290	1600/>5000	0.7/0.5 ^(g)
JWST ^(e)	K_{F200W}	27.8	-8.1	-2.0	310	1600/>5000	0.7/0.5 ^(g)
ELT ^(f) (Dec $\leq 0^\circ$)	<i>K</i>	28.5	-8.1	-2.0	400	750/>4000	1.0/0.7 ^(g)

Coverage	Passband	5 σ Depth (mag)	Peak GCLF (AB mag)	Peak offset & galaxy bk (mag)	d_{Max} (Mpc)	Bright candidates
~15 k sq.deg. ($ b \geq 23^\circ$)	VIS/H _E	~26.0	-8.3	-0.5 (-0.5)	50	400