


Einstein Telescope status and prospects

Some results of the PRIN project:


Multimessenger astronomy in the Einstein Telescope Era (METE)




M. Branchesi

Gran Sasso Science Institute

INFN/INAF/ASI




Einstein Telescope: the European next generation GW observatory



**Triangular (10 km arms)
or 2L shape (15 km arms) misaligned**

Underground
Cryogenic
Xylophone

...



EINSTEIN
TELESCOPE

ESFRI ROADMAP 2021

ET collaboration around 2000 members

Science design comparison in
Branchesi, Maggiore et al. 2024 JCAP

MAIN RESULTS:


- For compact binary coalescences, the 2L enables gains of at least a factor of 2–3 in both detection rates and the accuracy of parameter reconstruction with respect to ET-triangle
- This improvement is critical for exploratory science: early Universe, IMBHs, high-SNR “golden events” for precision tests of gravity, very well-localized sources
- For several science cases (including joint GW/GRB observations) the 2L only on the high-frequency IFOs allows to reach goals that would require high- and low-frequency IFOs in ET-triangle.

ETO+ETC: geometry comparison

science+risk, cost, governance, socio-economic impact



EINSTEIN
TELESCOPE



Three candidate sites
candidates to host ET:
EMR, Sardegna, and Lusastia



Italy and the Netherlands have pledged
to cover nearly half of the cost if their
proposed sites are selected.

Recently, Sardinia and Saxony have
signed a declaration of intents to
strengthen scientific collaboration and
to support the “2L” solution




Geometry and site selection expected by 2027

Operations by 2040

EXPECTED SENSITIVITY



ET Science in a nutshell



ET science book

Combination of:

- Larger distances
- Increased number of detections
- Detections with very high SNR

ET Science in a nutshell





ASTROPHYSICS

- **Black hole properties**
 - origin (stellar vs. primordial)
 - evolution, demography
- **Neutron star properties**
 - interior structure (QCD at ultra-high densities, exotic states of matter)
 - demography
- **Multi-band and -messenger astronomy**
 - joint GW/EM observations (GRB, kilonova,...)
 - multiband GW detection (LISA)
 - neutrinos
- **Detection of new astrophysical sources**
 - core collapse supernovae
 - isolated neutron stars
 - stochastic background of astrophysical origin

FUNDAMENTAL PHYSICS AND COSMOLOGY


- **The nature of compact objects**
 - near-horizon physics
 - tests of no-hair theorem
 - exotic compact objects
- **Tests of General Relativity**
 - post-Newtonian expansion
 - strong field regime
- **Dark matter**
 - primordial BHs
 - axion clouds, dark matter accreting on compact objects
- **Dark energy and modifications of gravity on cosmological scales**
 - dark energy equation of state
 - modified GW propagation
- **Stochastic backgrounds of cosmological origin**
 - inflation, phase transitions, cosmic strings

Multi-messenger in the ET era



Forecasting detection and parameter estimation capabilities

CBC populations




Dupletsa et al. 2023 A&C
Dupletsa et al. 2025 PRD

<https://github.com/janosch314/GWFish>
PRIN METE
PERIODIC ACME TUTORIAL

Python Fisher-Matrix
code/package

+GW waveforms

- It implements SNR and horizon estimate for SNe
- Beyond long-wavelength approximation for kHz studies (see for example Arnab et al. arXiv:2507.14071 on direct measurement of the accretion disk mass)
- It is use for ET, LGWA and multi-band studies (see for example Benetti et al. 2026, A&A for WD-WD)




COMPACT OBJECT BINARY POPULATIONS




EINSTEIN
TELESCOPE

BINARY NEUTRON-STAR MERGERS





BINARY BLACK-HOLE MERGERS



Sampling astrophysical populations
of binary system of compact objects
along the cosmic history of the
Universe

10^5 BNS detections per year
 10^5 BBH detections per year


ET sky-localization capabilities



ET low frequency sensitivity make it possible
To localize BNS!



- O(100) detections per year with sky-localization (90% c.r.) $< 100 \text{ sq. deg}$
- Early warning alerts!

Next generation GW effort worldwide



Cosmic Explorer: L shaped detectors, two sites
(40km, 20 km [option])

Network sky-localization capabilities




- $O(1000)$ detections per year with sky-localization (90% c.r.) $< 10 \text{ sq. deg}$


Branchesi et al. 2023, Dupletsa et al. 2023,
Ronchini et al. 2022



Pre-merger detections



Banerjee et al. 2023, A&A
Branchesi et al. 2023, JCAP



Five minutes before the merger:

- ET → O(10) detections with sky-localization < 30 deg²
- ET+CE → O(100) detections with sky-localization < 30 deg²

Getting closer to the merger time:

- improvement in sky localization capabilities
- more distant events can be localized

MM change of paradigm





Image Credit: Ghirlanda

RELATIVISTIC JET PHYSICS,
GRB EMISSION MECHANISMS,
COSMOLOGY and MODIFIED GRAVITY



KILONOVA PHYSICS,
NUCLEOSYNTHESIS,
NUCLEAR PHYSICS
and H0 ESTIMATE




HIGH-ENERGY

RELATIVISTIC JET PHYSICS,
GRB EMISSION MECHANISMS,
COSMOLOGY and MODIFIED GRAVITY

COSMOLOGY and MODIFIED GRAVITY


GW/Short GRB prompt emission




ET science book

Almost all detected short GRBs will have a GW counterpart

See Ronchini, MB et al. A&A 2022
Colombo et al. A&A 2025



AFTERGLOW EMISSION



ET science book





Image Credit: Ronchini




GW + γ -ray joint detections per year

SURVEY MODE

Fermi-GBM+ET



Fermi-GBM+(ET+CE)




Depending on the satellites, we will have **tens (e.g. THESEUS) to hundreds (e.g. HERMES)** of detections per year

Crucial instruments able to localize at arcmin-arcsec level to drive the ground-based follow-up

AFTERGLOW: Wide FoV X-ray telescopes (EP, THESEUS-SXI) increases the number of joint detections including off-axis GRB afterglow

Sensitive X-ray instruments such as **newATHENA** are required for the characterization

Ronchini, MB et al. A&A 2022



Implementation of the theoretical framework developed by Ronchini et al. 2022 used for THESEUS and GRINTA proposals

A new fast code almost ready to be released


MAGGPY




Provided the BNS GW observations, is there a tension among BNS merger populations and short GRBs?


Constraining BNS populations using Short GRB Observations+

De Santis, Ronchini, Santoliquido, Branchesi 2026 ready to be submitted to A&A






THESEUS and ET




GW observatories	Prompt
ET Δ 10 km	27 – 33
ET 2L 15 km	34 – 38
ET 2L 15 km + CE L 40 km	43 – 44

Expected number of well-localized THESEUS short GRBs with a joint GW detection over the nominal mission science time



Viewing angle vs. redshift distributions of THESEUS prompt and “orphan” afterglow detections



Afterglow-only cases up to viewing angle of ~14 deg are limited to $z < 1 \rightarrow$
3-4 other detections
Unprecedented precise parameter estimation from the GW


Post-O5 scenario 3A#
4-6 joint THESEUS+GW detections

Cosmology with joint observations of GWs and GRBs

- Mock MM data based on short GRB observed by Swift and Fermi
- Fisher matrix + agnostic reconstruction of the DE phenomenology (ML)





See *Cozzumbo et al. JCAP 2025*




HIGHLIGHTS:

- in combination with forthcoming cosmological surveys, around **40 GW-GRB detections** → unprecedented precision on H_0 and Ω_m , and DE density evolution
- required observatories capable of detecting GRBs and identifying their redshift.


CTAO and GW DETECTOR synergies



Pre-merger detections



Sky-localization capability:




Events per year

Detector	Ω [deg ²]	All orientations			
		15 min	5 min	1 min	0 min
ET + CE	100	442	1325	5075	123303
ET	100	90	130	208	436

Detector	Ω [deg ²]	Viewing angle ($<10^0$)			
		15 min	5 min	1 min	0 min
ET + CE	100	21	71	314	3376
ET	100	3	6	13	40


Observation strategy:



Follow-up pre-merger alerts with CTA

- Follow-up well localized sources ($< \text{FoV}$)
- Single shot observation
- Mosaic strategy
- Divergent pointing

Observation strategy: MST – one shot




ET+CE: around ten VHE counterparts can potentially be detected using 10% of the CTA time

THERMAL EMISSION - KILONOVAE

KILONOVA PHYSICS,
NUCLEOSYNTHESIS, NUCLEAR
PHYSICS and COSMOLOGY


PHYSICS and COSMOLOGY

Kilonova emission



ET science book


See Loffredo, Hazra, Dupletsa, MB et al. 2025, A&A
Colombo et al. A&A 2025



Einstein Telescope and Vera Rubin Observatory

64 simulations (release in zenodo Dupletsa et al. 2024)

- Two BNS merger populations (10 years)
- NS mass distribution: Gaussian and uniform
- EoSs: BLh and APR4
- Several network of detectors




Population KN light-curves

- Multi-component ejecta: dynamical + spiral wind + secular
 - KNe ejecta through numerical-relativity informed fits
 - Calibrated for **GW170817** and **GW190425**
 - Including **prompt collapse** of the remnant to black hole
- + sGRB afterglow

Realistic Observational strategy



- multi-epochs (2 nights)
- multi-filters (i and g)
- ToO observations, exposure = 600 s
- mosaic to cover the GW sky-loc

Gaussian NS mass distribution





Einstein Telescope and Vera Rubin Observatory

BLh uniform



BLh gaussian





- *multi-epochs (2 nights)*
- *multi-filters (i and g)*
- *ToO observations, exposure = 600 s*
- *mosaic to cover the GW sky-loc*



- ET → Rubin 10 –100 KNe per year
- ET-LVKI → Rubin detection increase by an order of magnitude wrt ET
- ET-CE → Rubin detection increase by 3 wrt ET+LVKI (not correspond to the increase of well-localised events because KN limited by Rubin detection efficiency)

Einstein Telescope and wide-field, high-multiplex spectroscopic facilities (WST)


WST-MOS Detectable KN as a function of time for ET 2L




Bisero et al. A&A 2026

BNS detected by ET in 10 yrs up to $z=1$ with sky-loc < 100 deg²

KN-magnitude at 12 hrs after the merger



Redshift distribution





Einstein Telescope and wide-field, high-multiplex spectroscopic facilities (WST)

Two strategies:

- WST alone performing a galaxy targeted research (peak of the probability regions with the IFS and the remaining area with MOS fibres) The second strategy that we consider
- WST in synergy with optical-NIR photometric observations - VRO

Number of galaxies in the comoving error volume of BNS



Bisero et al. A&A 2026

Golden events for WST follow-up:

- BNS with redshifts $z < 0.3$ and that have sky loc. $< 10 \text{ deg}^2$
- WST able to cover all the galaxies in the error volume with a limited number of exposures
- ~ 10 (ET-alone) to hundreds per year (ET+CE)







SUMMARY

In the ET era MM astronomy will become a routine

- Kilonovae will be detectable up to $z = 0.3-0.4$
[see Loffredo et al. 2025, Bisero et al. 2026, Colombo et al. 2025
Branchesi et al. 2024, Abac et al. 2026]
- GRBs will be the only counterparts detectable at high z
- Almost each detected GRB will have a GW counterpart
GRB localization will be crucial to drive the follow-up by ground
[see Ronchini et al. 2022, Colombo et al. 2025, Branchesi et al.
2024, Abac et al. 2026]

Early warning alerts critical to detect the prompt/early emission
[see Banjee et al. 2022, Branchesi et al. 2024, Abac et al. 2026]



Presented papers and coded were developed within the PRIN METE