

Magnetic fields in Galaxy clusters

Annalisa Bonafede

Università di Bologna

&

IRA INAF



THIS TALK

Part I:
techniques to constrain magnetic fields using
radio polarisation observations

Quiz 1



Quiz 2

Part II:
Magnetic field estimates in galaxy clusters
results, limitations and perspectives

Part III:
Origin and amplification of the
magnetic field in galaxy clusters






RADIO DIFFUSE EMISSION IN CLUSTERS



Radio halos, mini halos, and relics
probe the existence of magnetic fields but do not
offer a way to constrain them

EVIDENCE FOR B IN CLUSTERS: FILAMENTS AND SUBSTRUCTURES




HOW CAN WE CONSTRAIN MAGNETIC FIELDS IN CLUSTERS?


Back to the basics:

Polarisation of electromagnetic waves indicates the direction of oscillation of the **electric field** of the wave

Synchrotron emission is **linearly polarised**, with $\vec{E}$ perpendicular to $\vec{B}$ of the source



THE FARADAY EFFECT



$$n_{R,L} = \left(1 - \frac{\omega_p^2}{\omega^2 \pm \omega \Omega_e} \right)$$



ω_p = plasma frequency

ω = freq of emitted radiation

Ω_e = Lorentz freq of electrons



THE FARADAY EFFECT



$$\psi_{obs} = \psi_{int} + RM\lambda_{obs}^2 \quad RM = K \int_{los} B_{//} n_e dl$$

THE FARADAY EFFECT

$$\psi_{obs} = \psi_{int} + \text{RM} \lambda_{obs}^2 \qquad \text{RM} = 812 \int_{los} \frac{B_{//}}{\mu\text{G}} \frac{n_e}{\text{cm}^{-3}} \frac{dl}{\text{kpc}}$$

If we observe a **synchrotron source** in the background of a cluster at **multiple wavelengths**, we can derive the RM which gives us information about the magnetic field

- It's a **rotation** of the polarisation vector $\rightarrow$ the intensity of the polarisation is not changed
- The same RM will produce a **different** $\Delta\psi = \psi_{obs} - \psi_{int} \rightarrow$ large rotation at long λ



THE FARADAY EFFECT: EXAMPLE

$$\psi_{obs} = \psi_{int} + \text{RM} \lambda_{obs}^2 \qquad \text{RM} = 812 \int_{los} \frac{B_{//}}{\mu\text{G}} \frac{n_e}{\text{cm}^{-3}} \frac{dl}{\text{kpc}}$$

- A magnetic field of $1 \mu\text{G}$, in a density medium of 1 cm^{-3} , crossing $los=100 \text{ kpc}$ will produce
 $\text{RM} = 81.2 \text{ rad/m}^2$
- We measure angles $\Delta\psi = \text{RM} \lambda^2$ with errors! :
 - if we observe at 10 GHz $\rightarrow 0.073 \text{ radians} \sim 4^\circ$
 - if we observe at 2 GHz $\rightarrow 1.82 \text{ radians} \sim 104^\circ$
- If you expect low $\Delta\psi$, you need to observe at low frequencies



POLARIZATION: COMPLEX NOTATION


$$\vec{P} = p \cdot e^{i2(\psi_0 + RM\lambda^2)} = \underbrace{p \cdot e^{i2\psi_0}}_{\text{Intrinsic polarisation}} \underbrace{e^{2iRM\lambda^2}}_{\text{Faraday rotation just adding a phase term}}$$

Intrinsic polarisation

Faraday rotation
just adding a phase term


Obtaining the RM in practice:

- Observe at **several** λ a source in the background of a medium with B and free electrons
- Derive ψ_{obs} at different λ
- Fit ψ_{obs} versus λ^2 , the slope is the **RM**
- $RM > 0$ if the magnetic field is directed towards the observer along the line of sight



FROM FARADAY ROTATION TO B

- Every medium between you and the source will contribute to the RM if it contains B and free electrons (including our Galaxy, the Earth ionosphere, ...) → the resulting RM will be the sum of the different media




$$RM \propto \int_{Galaxy} B_{||} n_e dl + \int_{cluster} B_{||} n_e dl$$



FROM FARADAY ROTATION TO B

- The magnetic field and the gas density are not uniform: different regions of the background source will produce different RM




$$\langle RM \rangle \approx 0 \text{ but } \sigma_{RM} \neq 0,$$

- σ_{RM} is a better proxy for the magnetic field in the cluster
- The polarization intensity will decrease (depolarisation)

FROM FARADAY ROTATION TO B

In more complex situations, the rotation medium (cluster) may emit synchrotron radiation itself → different layer of the medium will produce different RM along the same line of sight



In these more complex, but realistic, cases, the polarisation angle is not going linear with λ^2

more sophisticated techniques have been developed



THE RM SYNTHESIS TECHNIQUE - ADVANCED -

Faraday depth $\Phi = \frac{d\psi_{obs}}{d\lambda^2}$

Observable $P(\lambda^2) = \int_{-\infty}^{+\infty} F(\Phi) \cdot e^{i2\Phi\lambda^2} d\Phi$ Contains the different RM along the line of sight

$P(\lambda^2) \longleftrightarrow F(\Phi)$ Fourier conjugates!

With the RM synthesis technique one can recover the Faraday rotation of the different emitting sources along the line of sight

This is the technique used nowadays



FARADAY RM: TAKE HOME MESSAGES

From multifrequency radio observations in polarisation


$$RM \propto \int B_{||} n_e dl$$

Magnetic field
along the los

Gas density
from X-ray
observations

RM is the observable!
Not B

If n_e and $\vec{B}$ are not uniform



$$\langle RM \rangle = 0$$

$$\sigma_{RM} \neq 0$$

σ_{RM} is a better proxy
for $\vec{B}$ than the RM

Regions with $\sigma_{RM} \neq 0$
will also decrease the
intensity of the
polarised radiation



Time for the first quiz



THIS TALK

Part I:
techniques to constrain magnetic fields using
radio polarisation observations


Part II:
**Magnetic field estimates in galaxy clusters
results, limitations and perspectives**

Part III:
Origin and amplification of the
magnetic field in galaxy clusters



RM in CLUSTERS: OBSERVATIONS

- Increase in RM value at low projected distances from the cluster center
- RM are positive and negative
- RM approaches 0 at $r \approx R_{vir}$




Bohringer et al (2016)



RM in CLUSTERS:OBSERVATIONS

- 124 clusters, for a total of 819 sources
- RM are positive and negative
- Note the higher values with respect to the plot before, due to the higher resolution and sensitivity






Osinga et al (2025)



RM MAPS of sources in CLUSTERS



- RM positive and negative
- $\sigma_{RM} \neq 0$
indicating a turbulent magnetic field



Credits: Bonafede, Vacca, Spasic

FROM RM IMAGES to MAGNETIC FIELDS ESTIMATE

Example:
The Coma cluster





FROM RM IMAGES to MAGNETIC FIELDS ESTIMATE

- Decreasing trend of $|\langle RM \rangle|$ versus distance from cluster centre
- Zero-order estimate:
 $n_e = 10^{-4} \text{ cm}^{-3}$
 $L = 1.5 \text{ Mpc}$
 RM of the central source: $RM = 250 \text{ rad/m}^2$

$$RM = 812 \int_{\text{los}} \frac{n_e}{\text{cm}^{-3}} \frac{B_{\parallel}}{\mu\text{G}} \frac{dl}{\text{kpc}}$$

$$B \approx \frac{RM}{812 n_e L} \approx 2 \mu\text{G}$$

- We are assuming:
 - constant B
 - constant n_e
 we know they're not! (RM varies)



RM image of
the central
source

FROM RM IMAGES to MAGNETIC FIELDS ESTIMATE

- Numerical approach (Murgia et al 2004, others followed - Bonafede- Osinga - ...)

$$RM = K \int_{los} n_e B_{||} dl$$

$$n_e(r) = n_e(0) \left[1 + \left(\frac{r}{r_c} \right)^2 \right]^{-3/2\beta}$$

β model, parameters
known from **X-ray**
observations

Let's model the magnetic field profile:

Assume that the magnetic field has
Gaussian random components that follow
a power spectrum

$$|B_{\Lambda}| \propto \Lambda^n$$

and a global profile

$$B(r) = B_0 \left(\frac{n_e}{n_0} \right)^{\eta}$$




FROM RM IMAGES to MAGNETIC FIELDS ESTIMATE


$$RM \propto \int n_e B_{||} dl$$

3D Numerical model for B, including a power spectrum and a scaling with gas density

$$B(r) = B_0 \left(\frac{n_e(r)}{n_e(0)} \right)^\eta$$



Simulated RM image




FROM RM IMAGES to MAGNETIC FIELDS ESTIMATE

Example:
The Coma cluster


$$B(r) = B_0 \left(\frac{n_e}{n_0} \right)^\eta$$
$$B \propto k^n; k_{min} < k < k_{max}$$

Free parameters:
 $B_0, \eta, k_{min}, k_{max}, n$

χ^2 plot



best fit profile



$B_0 = 4.7 \mu G, \eta = 0.5$
assuming a kolmogorov power
spectrum



MAGNETIC FIELD in the COMA CLUSTER


$$B_0 = 4.7\mu G, \eta = 0.5$$

$$B(r) = B_0 \left(\frac{n_e}{n_0} \right)^\eta$$

- $B \propto n^{0.5} \rightarrow$ magnetic energy density scaling like the thermal energy density

Note: flux-freezing would predict $B \propto n_e^{2/3}$

- Is magnetic field energetically important?
remember: β plasma parameter $\beta = \frac{n_e k_B T}{B^2 / 8\pi}$




Given $n_e = 10^{-3} cm^{-3}$, $T \sim 10^8 K$ $\beta \sim 300$
magnetic pressure subdominant with respect to
thermal pressure



MAGNETIC FIELD in OTHER CLUSTERS

In total, we have B estimates through
RM in 17 clusters

B of few -10 μG in the centres
scaling as $B \propto n_e^\eta$
with $\eta = 0.5 - 1$




Credits: Loi et al 2025



WHY SO FEW CLUSTERS WITH A CONSTRAINED B?

- Main limit: number of sources detectable in polarization
- Deep MeerKAT observations of the Fornax cluster:
80 sources / sq degree
(noise 3 μ Jy/beam, beam 13")
- obtained with 50 different pointings of 5h each




Credits: Loi et al 2025



WHY SO FEW CLUSTERS WITH A CONSTRAINED B?

How many sources in polarisation
can we detect with 5h
observing time?
(we are reaching the confusion
noise in total intensity)




Credits: Bonafede



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

MAGNETIC FIELDS FROM RM: CHALLENGES AND LIMITS

- Low number of **polarised** sources per cluster
- **Position** of the source along the line of sight unknown
- There could be a **local contribution** of RM for cluster members if for example due to local compression of the gas





MAGNETIC FIELDS FROM RM: PERSPECTIVES

Statistical study of
sample of clusters
with similar properties

Attempt by Osinga et
al (2024)
124 clusters with
known X-ray
properties

Clusters span a range
of z and mass too
large be considered
together




MAGNETIC FIELDS FROM RM: PERSPECTIVES with SKA

SKA mid
Band 2: 0.96 - 1.7 GHz

- N density of sources in polarisation:
- 15 min (mosaic) **110 sources/sq degree**
 - 10h (deep targeted field) **414 sources/sq degree**

A Coma like cluster



B in CLUSTERS: TAKE HOME MESSAGES

- Few clusters with reliable estimates of B through RM studies

- B modelled as
$$B(r) = B_0 \left(\frac{n_e(r)}{n_e(0)} \right)^\eta$$
$$B_0 \sim 1 - 10 \mu\text{G}$$
$$\eta \sim 0.5 - 1$$

- These values indicate a magnetic field which is dynamically subdominant, but relevant for small-scale processes (particle acceleration, thermal conduction, ...)



Time for the second quiz

THIS TALK

Part I:
techniques to constrain magnetic fields using
radio polarisation observations

Part II:
Magnetic field estimates in galaxy clusters
results, limitations and perspectives

Part III:
Origin and amplification of the
magnetic field in galaxy clusters



ORIGIN OF MAGNETIC FIELDS IN CLUSTERS

- We need to explain magnetic fields of 1-10 μG on Mpc scale
- No coherent circular motions (like spiral galaxies) $\rightarrow$ mechanisms like $\alpha - \Omega$ dynamo do not work
- Is it enough assume the same seed field needed to explain the magnetic field in galaxies?



ORIGIN OF MAGNETIC FIELDS IN CLUSTERS

- We need to explain magnetic fields of 1-10 μG on Mpc scale
- No coherent circular motions (like spiral galaxies) $\rightarrow$ mechanisms like $\alpha - \Omega$ dynamo do not work
- Is it enough assume the same seed field needed to explain the magnetic field in galaxies?

$$Br^2 = \text{const} \quad \text{from flux freezing}$$

$$B_{gal} r_{gal}^2 = B_{cluster} r_{cluster}^2$$

$$B_{gal} \approx 5\mu\text{G} \quad r_{gal} \approx 10 \text{ kpc} \quad r_{cluster} \approx 500 \text{ kpc}$$

$$B_{clusters} \approx 0.002\mu\text{G}$$


- The magnetic field must have been amplified



ORIGIN OF MAGNETIC FIELDS IN CLUSTERS

Fig. 1 Magnetic field strength as a function of over density in cosmological SPH simulations. Starting from 3 different cosmological seed field strengths: 2×10^{-13} G (dark green), 2×10^{-12} G (black), 8×10^{-12} G (green) (Dolag et al. 2008, 2005a). Adiabatic evolution solely by compression in grey. Runs with galactic seeds are in red and blue

Evolution of B
starting from
different seed
fields





Donnert et al (2019)




B AMPLIFICATION THROUGH SMALL-SCALE DYNAMO

- Fluid perturbation will induce **turbulence** in the intra-cluster medium, Turbulence decays to smaller and smaller scales
- In brief, **Kolmogorov theory of turbulence**: random fluid motion with velocity dispersion v on scale l (eddies) breaks up into two smaller eddies of size $l/2$ due to the convective term $\vec{v} \cdot \nabla \vec{v}$ in the fluid equation
→ Energy cascade



MAGNETIC FIELDS AMPLIFICATION IN CLUSTERS

- Energy cascade from large spatial scales (small k) to small partial scales (large k)



large spatial scale ————— small spatial scale

At some point, the scale reached by the eddies has a kinetic energy comparable to **viscous** forces or to the **Lorentz force**

At this scale, the kinetic energy of the flow is dissipated into thermal energy

The **magnetic field** is **stretched and folded** by the random velocity field

STRETCHING AND FOLDING OF MAGNETIC FIELD LINES: AMPLIFICATION

Credits:
Steinwandel et al
(2024)





Figure 1. Schematic sketch of the small-scale turbulent dynamo that illustrates the stretching, twisting, and folding of field lines driven by ICM turbulence. As magnetic tension becomes stronger due to subsequent stretching, twisting, and folding, the process slows down and finally saturates when the turbulent kinetic energy in the smallest eddies that drive the process are in equipartition with the magnetic energy density.





Exponential increase of the magnetic energy that generates an
“inverse cascade” from small to large scales

REMEMBER: flux freezing - incompressible fluid
any motion that increases l will also increase B : $B \propto l$

implies that: Field B_1 on scale l_1 stretched to scale l_2 : $B_2 = B_1 \frac{l_2}{l_1}$



B AMPLIFICATION THROUGH SMALL-SCALE DYNAMO




Exponential growth of the small scale dynamo does not proceed to infinity, as magnetic field starts to back react to the flow
“inverse cascade” up to a maximum scale l_B where B energy $\sim$ turbulent kinetic energy



B AMPLIFICATION THROUGH SMALL-SCALE DYNAMO

- The exponential growth of magnetic field in clusters is very fast → the dynamo saturates at the very beginning of cluster formation
- Every weak seed will be amplified by the dynamo




Dynamo will erase all imprints of the initial seed field



B AMPLIFICATION in CLUSTERS

- Dynamo is not the only process that shapes B in clusters:
 - clusters undergo **multiple mergers** and turbulent cascades start at different scales at once
 - Also **compression from shocks and shear motion** impact the shape and strength of the magnetic field




The resulting amplification of the magnetic field strength is a complex **interplay between compression and the small-scale dynamo**



RECENT UNEXPECTED RESULTS


- Radio halos found in high- z clusters
 $0.7 < z < 0.9$
Inverse Compton losses should be strong
 $\left(\frac{dE}{dt} \propto (1+z)^4\right)$
- To be visible in radio, B must be already amplified at $z \sim 0.7$ as in local clusters
- We need a fast amplifications (only few Gyrs from cluster formation)

Challenging dynamo models



ORIGIN AND AMPLIFICATION of B in CLUSTERS: TAKE HOME MESSAGES

- Magnetic field must be **amplified** to reach the μG level that we observe in clusters.
- The amplification depends on complex small-scale physics, likely involving **small-scale dynamos**
- The amplification must be fast! How much? It depends on the saturation value which is not constrained yet



we need robust **constrain on magnetic fields** on large samples of clusters to understand the amplification process



