

## Exceptional ALMA look at the anatomy of a luminous quasar host

*Thursday, 11 October 2018 11:45 (15 minutes)*

The first ALMA observation of the nearby quasar PDS 456 ( $z=0.184$ ) was designed to obtain the highest resolution map ever taken of the molecular gas in a hyper-luminous quasar. With a bolometric luminosity of  $2 \times 10^{47}$  erg/s ( $\sim$  Eddington luminosity) it can be regarded as the local counterpart of the quasars shining at  $z \sim 2$ , i.e. the peak of quasar luminosity density.

We are able to reveal in unprecedented detail (i.e.,  $0.2 \text{ arcsec} \sim 600 \text{ pc}$ ) the distribution of the molecular gas around a very powerful AGN and probe the relationship between nuclear and host galaxy properties at the brightest end of the luminosity function.

The most noticeable result is the discovery of a molecular outflow in PDS 456, which also shows the undisputedly most powerful, persistent, X-ray ultra-fast ( $0.25c$ ) wind discovered so far. This allows us to get new insights on how multi-phase AGN-driven outflows expand outwards. Interestingly, the CO outflow exhibits a very complex, unusual morphology with a compact, sub-kpc component and a  $\sim 3 \text{ kpc}$  extended, very diffuse approaching one. Finally, multiple gas-rich companions are visible around  $\sim 20 \text{ kpc}$  from PDS 456 supporting the merger-driven scenario for high-luminosity quasars.

### Affiliation

INAF-OAR

**Primary author:** PICONCELLI, Enrico (INAF - OARoma)

**Presenter:** PICONCELLI, Enrico (INAF - OARoma)

**Session Classification:** Outflows and Feedback processes