Contribution ID: 83

## Quasars as high redshift standard candles: the Lx-Luv relation at high redshift

Thursday, 11 October 2018 17:00 (15 minutes)

A tight non-linear relation between the X-ray and the optical-ultraviolet luminosity has been observed in AGN over 5 orders of magnitude and up to high redshift.

This suggests a coupling between the disk, emitting the primary radiation in the UV band, and the hot corona emitting in the X-ray.

In this work, we have studied the  $L_X - L_{UV}$  relation for a sample of high-redshift (z > 4) quasars, selected on the basis of their spectral properties and the quality of the available observations in both X-ray and optical/UV bands.

The relation shows no-evidence of evolution with redshift, which indicates a universal physical mechanism regulating the energy transfer process in the inner regions, and the observed dispersion is lower than in previous estimates.

The non-linearity of the relation provides a new, powerful way to estimate the absolute luminosity, turning quasars into a new class of standard candles that can provide an important contribution in the determination of the cosmological parameters, probing cosmological time not achievable with other observational methods.

## Affiliation

Dipartimento di Fisica e Astronomia, Università di Bologna

**Primary authors:** SALVESTRINI, Francesco (DIFA- Univ. Bologna; INAF - OAS Bologna ); Prof. RISALITI, Guido (Università di Firenze)

Co-authors: Dr BISOGNI, Susanna (INAF - Arcetri); Dr LUSSO, Elisabeta (Durham University)

Presenter: SALVESTRINI, Francesco (DIFA- Univ. Bologna; INAF - OAS Bologna )

Session Classification: Cosmology and High-redshift