



Contribution ID: 44

Type: Talk

## X-ray variability plane using NLSy1 and Sy1: Importance of obscuration

Friday, 13 April 2018 10:30 (20 minutes)

**Scaling relations** are the most powerful astrophysical tools to set constraints to the physical mechanisms of astronomical sources and to infer properties for objects where they cannot be accessed directly. We have re-investigated one of these scaling relations using NLSy1 and Sy1; the so-called **X-ray variability plane** (or mass-luminosity-timescale relation, McHardy et al. 2006). This relation links the power-spectral density (PSD) break frequency with the SMBH mass and the bolometric luminosity. We used all available XMM-Newton observations of a sample of 22 Sy1 and NLSy1 to study the PSD and spectra in short segments within each observation. This allows us to report for the first time that the PSD break frequency varies for each object, showing variations in 19 out of the 22 AGN analyzed. Our analysis of the variability plane confirms the relation between the break frequency and the SMBH mass and finds that the **obscuration along the line of sight** (or the variations on the obscuration using its standard deviation) **is also a required parameter**. We constrain a new variability plane of the form:  $\log(\nu_{\text{Break}}) = -A \log(M_{\text{BH}}) + B \log(N_{\text{H}}) - C$  (or  $\log(\nu_{\text{Break}}) = -A \log(M_{\text{BH}}) + B (\log(N_{\text{H}}) + C)$ ). The X-ray variability plane found by McHardy et al. (2006) is roughly recovered when we use unobscured segments. We speculate the PSD shape is related with the outflowing wind close to the accretion disk in NLSy1 and Sy1.

### Motivation

### Grant

no

**Primary author:** GONZALEZ-MARTIN, Omaira (IRyA)

**Presenter:** GONZALEZ-MARTIN, Omaira (IRyA)