



X-RAY ASTRONOMY 2019

Current Challenges and New Frontiers in the Next Decade

8-13 September 2019
CNR/INAF Research Area, Bologna, Italy

Contribution ID: 99

Type: **Contributed**

AGN radiative feedback: the effective Eddington limit for dusty gas

Wednesday, 11 September 2019 14:45 (15 minutes)

AGN radiative feedback, driven by radiation pressure on dust, is a key physical mechanism connecting the accreting black hole to its surrounding environment. The actual importance of radiative feedback can be observationally tested by analysing how X-ray-selected AGN samples populate the so-called “NH-lambda plane”, defined by the column density versus the Eddington ratio. A “forbidden” region occurs in this plane, where obscuring clouds cannot be long-lived, and should be outflowing. We show how the inclusion of radiation trapping leads to an enhanced forbidden region, such that even Compton-thick material can potentially be disrupted by sub-Eddington luminosities. Comparing our model results to the most complete sample of local AGNs with measured X-ray properties, we obtain a nice agreement. In addition, by properly including radiation trapping, we can account for the dynamics and energetics of AGN-driven outflows observed on galactic scales. I will discuss how such radiative feedback may regulate both the AGN obscuration and outflow properties in the context of black hole-host galaxy co-evolution.

Topic

Active Galactic Nuclei: accretion physics and evolution across cosmic time

Affiliation

University of Zurich

Primary author: Dr ISHIBASHI, Wako (University of Zurich)

Presenter: Dr ISHIBASHI, Wako (University of Zurich)

Session Classification: ACTIVE GALACTIC NUCLEI