



X-RAY ASTRONOMY 2019

Current Challenges and New Frontiers in the Next Decade

8-13 September 2019
CNR/INAF Research Area, Bologna, Italy

Contribution ID: 39

Type: **Contributed**

Evidence for radiation pressure compression in the X-ray narrow-line region of Seyfert galaxies

Wednesday, September 11, 2019 11:40 AM (15 minutes)

The observed overlap between soft X-ray emission and the NLR in obscured AGN is commonly interpreted as evidence for a constant gas pressure multiphase medium. Radiation pressure compression (RPC) also leads to a density distribution, since a gas pressure (hence density) gradient must arise within each cloud to counteract the ionizing radiation pressure. RPC leads to a well-defined ionization distribution, and a differential emission measure (DEM) distribution with a universal slope of -0.9 . In contrast, a multiphase medium does not predict the form of the DEM. The observed DEMs of obscured AGN with XMM-Newton RGS spectra (the CHRESOS sample) are in striking agreement with the predicted RPC DEM, providing a clear signature that RPC is the dominant mechanism in the X-ray NLR. In contrast with the constant gas pressure multiphase medium, RPC further predicts an increasing gas pressure with decreasing ionization, which can be tested with future X-ray missions using density diagnostics.

Topic

Active Galactic Nuclei: accretion physics and evolution across cosmic time

Affiliation

Università degli Studi Roma Tre

Author: Prof. BIANCHI, Stefano (Università degli Studi Roma Tre)

Presenter: Prof. BIANCHI, Stefano (Università degli Studi Roma Tre)

Session Classification: ACTIVE GALACTIC NUCLEI