



X-RAY ASTRONOMY 2019

Current Challenges and New Frontiers in the Next Decade

8-13 September 2019
CNR/INAF Research Area, Bologna, Italy

Contribution ID: 160

Type: **Contributed**

Tension with the Λ CDM model from a Hubble diagram of quasars.

Thursday, September 12, 2019 9:30 AM (15 minutes)

The cosmological concordance model (Λ CDM) well accounts for a wealth of observations, from the existence of Cosmic Microwave background (CMB) to the discovery of the accelerated expansion of the universe from Type Ia supernovae. Yet, it assumes a still unknown form of dark energy and matter and some tensions arose recently as, for instance, the discovery of a $>3\sigma$ discrepancy between the local (Riess et al. 2016) and Planck (Aghanim et al. 2016) measurement of H_0 . In addition, the Λ CDM model is poorly tested in the redshift interval between the farthest observed Type Ia supernovae ($z \sim 1.4$) and that of the CMB ($z \sim 1100$). We present new measurements of the expansion rate of the Universe in the redshift range $z=0.5-5.5$ based on a Hubble diagram of quasars. The distance of quasars have been estimated from the observed non-linear relation between the X-ray and ultraviolet emission, following a method developed by our group. The distance-redshift relation of quasars at $z < 1.4$ is in agreement with that of supernovae and with the concordance model. Nonetheless, a deviation from the Λ CDM model emerges at higher redshift, with a statistical significance of $\sim 4\sigma$. We found that, if an evolution of the dark energy equation of state is allowed, our data suggest a dark energy density increasing with time.

Topic

Affiliation

Dipartimento di Fisica e Astronomia, Università di Firenze

Author: LUSSO, Elisabeta (Dipartimento di Fisica e Astronomia, Università di Firenze)

Presenter: LUSSO, Elisabeta (Dipartimento di Fisica e Astronomia, Università di Firenze)

Session Classification: ACTIVE GALACTIC NUCLEI