X-RAY ASTRONOMY 2019



8-13 September 2019 CNR/INAF Research Area, Bologna, Italy

Contribution ID: 75 Type: Poster

The X-ray emission from hot subdwarf stars

Friday, 13 September 2019 15:13 (1 minute)

In the last decade the high throughput and spectral resolution of the *XMM-Newton* and *Chandra* X-ray telescopes has allowed us to investigate the X-ray emission from hot subdwarf stars. Up to now, we have detected in X-rays five sdO stars. All of them show intrinsic X-ray emission and their spectrum can be described with the sum of two or three thermal-plasma components, as in the case of normal O-type stars. Therefore, the X-ray emission seems to originate from shock instabilities in the wind. These results show that the X-ray observation of hot subdwarf stars is essential for understanding their properties.

Topic

Compact and diffuse sources in galaxies and in the Galactic Center

Affiliation

INAF - IASF Milano

Primary authors: LA PALOMBARA, Nicola (Istituto Nazionale di Astrofisica (INAF)); MEREGHETTI, Sandro (Istituto Nazionale di Astrofisica (INAF)); Dr ESPOSITO, Paolo (INAF-IASF Milano); Dr TIENGO, Andrea (IUSS Pavia)

Presenter: LA PALOMBARA, Nicola (Istituto Nazionale di Astrofisica (INAF))

Session Classification: POSTER SESSION