



Contribution ID: 59

Type: not specified

## Early Metal Enrichment and Multiphase Interstellar Media in a Galaxy at Redshift 8.312

Wednesday, 16 October 2019 09:25 (15 minutes)

Contributed talk

Abstract:

“How and when metal/dust enrichment happened in the epoch of reionization is one of the most fundamental questions in modern astronomy. Recent Planck results suggest an instantaneous reionization redshift of  $z = 7.7 \pm 0.8$ , and the latest Hubble Space Telescope surveys have revealed strong evolution of the ultraviolet luminosity function from  $z \sim 10$  to  $\sim 8$ , implying a rapid increase of the cosmic star-formation rate density within a very short time-scale ( $< 200$  Myr). Hence, characterizing star formation and interstellar medium of galaxies in the heart of the reionization era ( $z \sim 8$ ) is an important next step to our understanding of the earliest evolution of galaxies.

We present ALMA detections of the [OIII] 88  $\mu\text{m}$ , [CII] 158  $\mu\text{m}$  lines and 850  $\mu\text{m}$  dust continuum emission in a Y-dropout galaxy, MACS0416\_Y1. The [OIII] and [CII] detections unambiguously confirm its spectroscopic redshift to be  $z = 8.312$ , making this object one of the furthest galaxies ever identified spectroscopically. The ultraviolet-to-far infrared spectral energy distribution (SED) modeling where the [OIII] emissivity model is incorporated suggests the presence of a young (age  $\sim 4$  Myr), star-forming (SFR  $\sim 60$  Msun/yr), moderately metal-polluted ( $Z \sim 0.2 Z_{\text{sun}}$ ) stellar component. An analytic dust mass evolution model with a single episode of star formation does not, however, reproduce the metallicity and the inferred dust mass of  $4 \times 10^6$  Msun in the stellar age, suggesting an underlying evolved stellar component as the origin of the dust mass.

A high [OIII]-to-[CII] luminosity ratio ( $\sim 6$ ) implies that typical H II regions are radiation-bounded, which is consistent with the inferred zero escape fraction of Lyman continuum photons obtained in the SED fits. It also suggests that the bulk of photodissociation regions are ‘truncated’ and are not terminated by molecular gas, although this would contradict the presence of dust, which should be co-spatial with molecular gas. Thus, a possible explanation for the observed fact is patchy/porous geometry of dusty molecular clouds, photodissociation regions, and highly ionized gas on sub-kpc scales, which are all suggested by state-of-the-art hydrodynamic simulations of galaxy formation.”

**Presenter:** Dr TAMURA, Yoichi

**Session Classification:** Cosmology