



Contribution ID: 80

Type: not specified

Shaping of the stellar wind of evolved red giants and their successors by (sub-)stellar binary companions

Thursday, October 17, 2019 5:35 PM (15 minutes)

Contributed talk

Abstract:

Planetary nebulae (PNe) reveal a wide range of morphologies. Bipolarity is the main characteristic, but jets and tori are also detected. Several contending theories of the evolution from a (roughly) spherically symmetric Asymptotic Giant Branch (AGB) stellar wind to a very non-spherical PN have emerged. Here, we present the first high-spatial resolution observational campaign of a large sample of oxygen-rich AGB stellar winds obtained in the ALMA Large Program ATOMIUM ('ALMA Tracing the Origins of Molecules forming dUst in oxygen-rich M-type stars'). The data constitute the first observational proof that (sub-)stellar binary activity is the dominant shaping mechanism of AGB stellar winds, and their successors the PNe. We show that low mass-loss rate oxygen-rich AGB winds are more readily prone to complex structural deformations owing to their slow wind acceleration, whereas a binary-induced spiral structure is more prevalent in other classes of AGB stellar winds. These results resolve several previously unexplained phenomena —including the absence of detached shells around oxygen-rich AGB stars and disks around carbon-rich PNe —and have critical implications for the formation of type Ia supernovae.

Presenter: Dr DECIN, Leen

Session Classification: Stellar Evolution