Congresso Nazionale di Astrochimica e Astrobiologia (proto-) planetaria



Contribution ID: 14

Type: Oral

Deuterated amidogen in the laboratory and in Space

Monday 21 October 2019 12:20 (25 minutes)

Light nitrogen hydrides as imidogen (NH), amidogen (NH₂) and ammonia (NH₃) are located at the very beginning of the reaction network which leads to the formation of complex species, and are thus of primary importance to constrain the models describing nitrogen chemistry in the ISM. They are formed by a chain of hydrogen abstraction reactions starting from:

$$N^+ + H_2 \longrightarrow NH^+ + H$$
 , . . .

which also produces the ions $NH_2^+ NH_3^+$ and NH_4^+ , and then the corresponding neutral hydrides by dissociative recombination. If H_2 is replaced with its singly deuterated form HD, this ion–neutral process has a much reduced endothermicity and can thus drive deuterium into the molecular form, enhancing the abundance of ND, NHD, and the NH_nD_{3-n} isotopologues.

Amidogen (NH₂) has been detected in the interstellar medium: towards the massive star forming region SgrB2 [1], and in the low-mass proto-binary system IRAS 16293-2422 [2]. On the other hand, its deuterated variants NHD and ND₂ have not been identified in Space to date due to the incomplete knowledge of their rotational spectrum. Recent laboratory studies performed in Bologna [3] and at MPE Garching [4] have provided very accurate rest-frequencies for these isotopologues at sub-millimetre and THz regimes, thus allowing for sound searches for their spectral features in *Herschel*/HIFI archival data.

Accurately examining the spectral scan observations performed towards IRAS16293-2422 on the framework of the CHESS Herschel key project [5], we tentatively identified for the first time singly- and doubly-deuterated amidogen in an astrophysical source. The $1_{11} - 0_{00}$, J = 3/2 - 1/2 line of NHD and the $2_{12} - 1_{01}$, J = 5/2 - 3/2 line of ND₂ were detected in absorption towards the protostar sub-millimetre continuum. The hyperfine analysis of NHD and ND₂ spectral patterns, together with a reanalysis of the NH₂ spectrum, provided a measure of the amidogen deuteration in this proto-stellar environment.

References:

[1] E.F. van Dishoeck, D.J. Jansen, P. Schilke and T.G. Philips, ApJ, 416, L83 (1993)

[2] P. Hily-Blant, S. Maret, A. Bacmann, et al., A&A, 521, L52 (2010)

- [3] M. Melosso, C. Degli Esposti, L. Dore, *ApJS*, 233, 15 (2017)
- [4] L. Bizzocchi et al., manuscript in preparation
- [5] C. Ceccarelli, A. Bacmann, A. Boogert, et al., A&A, 521, L22 (2010)

Authors: BIZZOCCHI, Luca (Max-Planck-Institut für extraterrestrische Physik); GIULIANO, Barbara Michela (Max Planck Institute for Extraterrestrial Physics); MELOSSO, Mattia (Dipartimento di Chimica "G. Ciamician", Università di Bologna); Dr DORE, Luca (Dipartimento di Chimica "G. Ciamician", Università di Bologna); Dr TAMASSIA, Filippo (Dipartimento di Chimica Industriale "Toso Montanari", Università di Bologna); Prof. CASELLI, Paola

Presenter: BIZZOCCHI, Luca (Max-Planck-Institut für extraterrestrische Physik)

Session Classification: Protostellar chemistry