

Space Weather Group. University of Alcalá

# **Analysis of the solar wind distribution functions at 1 AU**

C. Larrodera, C. Cid  
carlos.larrodera@edu.uah.es

September 6, 2021



Introduction

Data

Bi-Gaussian approach

Solar cycle evolution

ICMEs Identification

Conclusions



- ▶ The solar wind is classified in slow and fast wind (bulk solar wind) and transient events
- ▶ Different distribution functions have been proposed to characterize the solar wind distribution, e.g. Burlaga and King (1979); Li et al. (2016)
- ▶ Solar wind magnitudes, like proton speed, evolve dynamically between Sun and Earth. Nevertheless, no major changes are expected in the composition
- ▶ Solar wind composition is used as a signature of interplanetary coronal mass ejection (Heidrich-Meisner et al., 2016)
- ▶ Average iron charge state ( $\langle Q_{Fe} \rangle$ ) values above 12 show the presence of ICMEs (Lepri et al., 2001; Lepri, 2004)



- ▶ We use data from the Advanced Composition Explorer (ACE) at the L1 point
- ▶ The data range is from 1998 to 2017
- ▶ The sources of the data are the instruments
  - ▶ Magnetic Field Experiment (MAG)
  - ▶ Solar Wind Ion Composition Spectrometer (SWICS)
  - ▶ Solar Wind Electron, Proton and Alpha Monitor (SWEPAM)




- ▶ We propose a bi-Gaussian distribution function to characterize the solar wind distribution

$$bG(x) = h_1 \cdot \exp\left(\frac{-(x - p_1)^2}{2w_1^2}\right) + h_2 \cdot \exp\left(\frac{-(x - p_2)^2}{2w_2^2}\right)$$

- ▶  $h$  is the height of the peak,  $p$  the position of the center and  $w$  the Gaussian RMS




- ▶ We have applied the bi-Gaussian distribution function to the whole data set of:
  - ▶ Proton speed
  - ▶ Proton temperature
  - ▶ Proton density
  - ▶ Magnetic field magnitude



**Figure 1:** Solar wind distribution for different magnitudes,  $n_p$ ,  $T_p$ ,  $V_p$  and  $B$  for the whole ACE data set (Larrodera and Cid, 2020a)



- We have also applied it to the whole data set of the average iron charge state  $\langle Q_{Fe} \rangle$




**Figure 2:** Solar wind distribution for  $\langle Q_{Fe} \rangle$  for the whole ACE data set (Larrodera and Cid, 2020b)


# Solar cycle evolution



- ▶ We have applied the bi-Gaussian approach also to the yearly data set
- ▶ We are able to study how the position of the peaks of the Gaussian PDF evolves
- ▶ We compare the position of the peaks with the Sunspot Number in order to study the correlation with the Solar Cycle



**Figure 3:** Scatter plots for  $\langle Q_{Fe} \rangle$  against the Sunspot Number. (Larrodera and Cid, 2020b)



**Figure 4:** Scatter plot of  $n_p$ ,  $T_p$ ,  $v_p$  and  $B$  against the sunspot number. (Larrodera and Cid, 2020a)





- ▶ Large deviation from typical values of  $\langle Q_{Fe} \rangle$  are related with ICMEs
- ▶ Considering  $\langle Q_{Fe} \rangle > 12$  at least for 10 hours we found 27 events:
  - ▶ 'Extended': Events where an extension of catalogued ICMEs will covered them
  - ▶ 'New': Events not previously catalogued



- ▶ The bi-Gaussian function properly reproduces the bulk solar wind
- ▶ The five magnitudes analyzed show a bimodal distribution
- ▶ These results suggest that the bulk solar wind at 1 AU is bi-modal
- ▶ Some fitting parameters show a strong correlation with the solar cycle
- ▶  $\langle Q_{Fe} \rangle$  allow us to locate ICMEs previously not cataloged
- ▶  $\langle Q_{Fe} \rangle > 12$  is a sufficient signature to identify ICMEs and its boundaries



- ▶ A complete description of our research can be found in the published papers:
  - ▶ <https://www.aanda.org/articles/aa/abs/2020/03/aa37307-19/aa37307-19.html>
  - ▶ <https://link.springer.com/article/10.1007/s11207-020-01727-8>
- ▶ For further questions, please contact me at:  
**carlos.larrodera@edu.uah.es**



- L. F. Burlaga and J. H. King. Intense interplanetary magnetic fields observed by geocentric spacecraft during 1963-1975. *Journal of Geophysical Research*, 84(A11):6633-6640, Nov 1979. doi: 10.1029/JA084iA11p06633.
- Verena Heidrich-Meisner, Thies Peleikis, Martin Kruse, Lars Berger, and Robert Wimmer-Schweingruber. Observations of high and low Fe charge states in individual solar wind streams with coronal-hole origin. *Astronomy & Astrophysics*, 593:A70, September 2016. doi: 10.1051/0004-6361/201527998.
- C. Larrodera and C. Cid. Bimodal distribution of the solar wind at 1 au. *Astronomy & Astrophysics*, 635:A44, 2020a.
- C. Larrodera and C. Cid. The distribution function of the average iron charge state at 1 au: from a bimodal wind to icme identification. *In prep*, 2020b.
- S. T. Lepri. Iron charge state distributions as an indicator of hot ICMEs: Possible sources and temporal and spatial variations during solar maximum. *Journal of Geophysical Research*, 109(A1), 2004. doi: 10.1029/2003ja009954.
- S. T. Lepri, T. H. Zurbuchen, L. A. Fisk, I. G. Richardson, H. V. Cane, and



## Introduction

- The solar wind is classified in slow and fast wind (bulk solar wind) and transient events
- Different distribution functions have been proposed to characterize the solar wind distribution, e.g. Burlaga and King (1979); Li et al. (2016)
- Solar wind magnitudes, like proton speed, evolve dynamically between Sun and Earth. Nevertheless, no major changes are expected in the composition
- Solar wind composition is used as a signature of interplanetary coronal mass ejection (Heidrich-Meisner et al., 2016)
- Average iron charge state ( $\langle Q_{Fe} \rangle$ ) values above 12 show the presence of ICMEs (Lepri et al., 2001; Lepri, 2004)

## Data set

- We use data from the Advanced Composition Explorer (ACE) at the L1 point
- The data range is from 1998 to 2017
- The sources of the data are the instruments
  - Magnetic Field Experiment (MAG)
  - Solar Wind Ion Composition Spectrometer (SWICS)
  - Solar Wind Electron, Proton and Alpha Monitor (SWEPAM)

## Bi-Gaussian approach

- We propose a bi-Gaussian distribution function to characterize the solar wind distribution

$$bG(x) = h_1 \cdot \exp\left(\frac{-(x - p_1)^2}{2w_1^2}\right) + h_2 \cdot \exp\left(\frac{-(x - p_2)^2}{2w_2^2}\right)$$

- $h$  is the height of the peak,  $p$  the position of the center and  $w$  the Gaussian RMS

## Bi-Gaussian approach. Dynamic magnitudes

- We have applied the bi-Gaussian distribution function to the whole data set of:
  - Proton speed
  - Proton temperature
  - Proton density
  - Magnetic field magnitude




Figure 1: Solar wind distribution for different magnitudes,  $n_p$ ,  $T_p$ ,  $V_p$  and  $B$  for the whole ACE data set (Larrodera and Cid, 2020a)

## Bi-Gaussian approach. Composition

- We have also applied it to the whole data set of the average iron charge state ( $\langle Q_{Fe} \rangle$ )




Figure 2: Solar wind distribution for  $\langle Q_{Fe} \rangle$  for the whole ACE data set (Larrodera and Cid, 2020b)

## Solar Cycle evolution

- We have applied the bi-Gaussian approach also to the yearly data set
- We are able to study how the position of the peaks of the Gaussian PDF evolves
- We compare the position of the peaks with the Sunspot Number in order to study the correlation with the Solar Cycle




Figure 3: Scatter plots for  $\langle Q_{Fe} \rangle$  against the sunspot number. (Larrodera and Cid, 2020b)




Figure 4: Scatter plot of  $n_p$ ,  $T_p$ ,  $v_p$  and  $B$  against the sunspot number. (Larrodera and Cid, 2020a)

## ICMEs identification

- Large deviation from typical values of  $\langle Q_{Fe} \rangle$  are related with ICMEs
- Considering  $\langle Q_{Fe} \rangle > 12$  at least for 10 hours we found 27 events:
  - ‘Extended’: Events where an extension of catalogued ICMEs will covered them
  - ‘New’: Events not previously catalogued

## Conclusions

- The bi-Gaussian function properly reproduces the bulk solar wind
- The five magnitudes analyzed show a bimodal distribution
- These results suggest that the bulk solar wind at 1 AU is bi-modal
- Some fitting parameters show a strong correlation with the solar cycle
- $\langle Q_{Fe} \rangle$  allow us to locate ICMEs previously not cataloged
- $\langle Q_{Fe} \rangle > 12$  is a sufficient signature to identify ICMEs and its boundaries

## Complete information

- A complete description of our research can be found in the published papers:
  - <https://www.aanda.org/articles/aa/abs/2020/03/aa37307-19/aa37307-19.html>
  - <https://link.springer.com/article/10.1007/s11207-020-01727-8>
- For further questions, please contact me at: [carlos.larrodera@edu.uah.es](mailto:carlos.larrodera@edu.uah.es)