

Contribution ID: 288

Type: Poster

Inference of log(gf) values for atomic lines in NUV spectral range

Monday, 6 September 2021 16:09 (13 minutes)

The third flight of the Sunrise observatory carries the new Sunrise UV Spectropolarimeter and Imager (SUSI) which will cover the spectral range from 310 nm to 410 nm. In this region, the number density of spectral lines is at least four times larger than in the visible/infrared part of the spectrum, including a significant number of chromospheric lines. To extract accurate physical parameters of the solar atmosphere from observations with SUSI we need precise atomic line parameters, e.g., the log (gf) value. Here we review an extension of the spectropolarimetric inversion scheme to account for coupled inversion of atomic line parameters. In this scheme, atomic line parameters are inverted using information from the whole field of view. Additionally, we discuss the potential of this inversion scheme using synthetic spectra computed from 3D realistic magnetohydrodynamic simulations.

Primary author: VUKADINOVIC, Dusan

Co-authors: NARAYANAMURTHY, Smitha (Max Planck Institute for Solar System Research); LAGG, Andreas (Max Planck Institute for Solar System Research); VAN NOORT, Michiel (Max Planck Institute for Solar System Research); SOLANKI, Sami (Max Planck Institute for Solar System Research)

Presenter: VUKADINOVIC, Dusan

Session Classification: Poster Session 2.2

Track Classification: Session 2 - The Solar Atmosphere: Heating, Dynamics and Coupling