

Contribution ID: 319

Type: Poster

## Hot X-ray Onsets of Solar Flares

Wednesday, 8 September 2021 15:18 (13 minutes)

The study of the localized plasma conditions before the impulsive phase of a solar flare can help us understand the physical processes that occur leading up to the main flare energy release. Here, we present evidence of a hot X-ray 'onset'interval of enhanced isothermal plasma temperatures in the range of 10-15 MK up to tens of seconds prior to the flare's impulsive phase. This 'hot onset'interval occurs during the pre-flare time during which elevated soft X-ray flux is detected, but prior to detectable hard X-ray emission. The isothermal temperatures, estimated by the Geostationary Operational Environmental Satellite (GOES) X-ray sensor, and confirmed with data from RHESSI, show no signs of gradual increase and occurs regardless of flare classification or configuration. In a small sample of four representative flare events, we identify this early hot onset soft X-ray emission mainly within footpoint and low-lying loops, rather than with coronal structures, based on images from the Atmospheric Imaging Assembly (AIA) and the use of limb occultation. These hot X-ray onsets appear before there is evidence of collisional heating by non-thermal electrons, and hence challenges the standard flare heating modelling techniques.

## Student poster?

Primary author: SIMÕES, Paulo (CRAAM/Mackenzie)

**Co-authors:** Dr HUDSON, Hugh (University of Glasgow/University of California Berkeley); Prof. FLETCHER, Lyndsay (University of Glasgow); HAYES, Laura (Dublin Institute of Advanced Studies (DIAS)); Dr HANNAH, Iain (University of Glasgow)

Presenter: SIMÕES, Paulo (CRAAM/Mackenzie)

Session Classification: Poster Session 7.5

**Track Classification:** Session 4 - From Radio to Gamma Rays: Near-Sun Manifestations and Triggering of Solar Flares and Coronal Mass Ejections