



Contribution ID: 131

Type: **Oral**

### **Jets in Blazars as seen by POLAMI: Polarization Monitoring of AGN at Millimeter Wavelengths with the IRAM 30m Telescope**

*Monday, 29 August 2022 10:15 (15 minutes)*

A Large Program at the IRAM 30m Telescope called POLAMI observes since 2007, with a time sampling of ~2 weeks, the four Stokes parameters of the 3.5 and 1.3 mm emission of a sample of ~40 of the brighter sources in the northern sky. This contribution outlines the most salient scientific results obtained from the first detailed analysis of the data on which regards to the structure and magnetic field of the innermost blazar jet regions. In particular, the analysis of the variability of our data implies that shorter mm emission should come from smaller regions with progressively better magnetic field order, one-zone models are definitely excluded by the general properties of mm polarization of blazars, blazar jets are not compatible with axisymmetric geometries in general on which regards to their polarization emission. Moreover, variable circular polarization emission seems to be present in most blazars at mm wavelengths at levels of ~2% or larger.

**Primary author:** Dr AGUDO, Ivan (Instituto de Astrofísica de Andalucía (CSIC))

**Presenter:** Dr AGUDO, Ivan (Instituto de Astrofísica de Andalucía (CSIC))

**Session Classification:** Polarization and AGN and blazars